

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C12N 5/04, 15/63, 15/82, 15/83, 15/90, G01N 33/556	A1	(11) International Publication Number: WO 00/46350 (43) International Publication Date: 10 August 2000 (10.08.00)
(21) International Application Number: PCT/US00/02971 (22) International Filing Date: 4 February 2000 (04.02.00) (30) Priority Data: 60/118,867 5 February 1999 (05.02.99) US (71) Applicant: THOMAS JEFFERSON UNIVERSITY [US/US]; 11th and Walnut Streets, Philadelphia, PA 19107 (US). (72) Inventors: KOPROWSKI, Hilary; 334 Fairhill Road, Wyn- newood, PA 19096 (US). YUSIBOV, Vidadi; 1808 Manor Road, Havertown, PA 19083 (US). (74) Agent: MCNICHOL, William, J., Jr.; Reed Smith Shaw & McClay LLP, 2500 One Liberty Place, Philadelphia, PA 19103-7301 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NI., PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: PRODUCTION OF BIOMEDICAL PEPTIDES AND PROTEINS IN PLANTS USING TRANSCOMPLEMENTATION SYSTEMS (57) Abstract <p>The present invention discloses novel methods for producing foreign polypeptides in a host plant using recombinant viruses through functional complementation systems. The methods involve constructing suitable recombinant viral vectors that are capable of systemic infection and infecting the host plants with one or more recombinant viral vectors. The methods are also directed to infecting a host plant that is transgenic for expressing replicase genes of a virus, wherein the transgenic plant expressing a virus replicase genes complements the virus replicase function. The invention also discloses compositions comprising recombinant viral vectors capable of systemic infection for producing foreign polypeptides in a host plant.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**PRODUCTION OF BIOMEDICAL PEPTIDES AND PROTEINS IN PLANTS
USING TRANSCOMPLEMENTATION SYSTEMS**

5

This application claims the priority of U.S. Provisional Application No. 60/118867, filed February 5, 1999.

FIELD OF THE INVENTION

10

The present invention is in the fields of genetic engineering and molecular farming, and especially provides methods for systemically producing foreign polypeptides and polynucleotides in plants using recombinant viral vectors and transcomplementation systems. Also provided are compositions for production of such polypeptides and polynucleotides in plants.

15

BACKGROUND OF THE INVENTION

In recent years, plants have been actively targeted for the production of medically important proteins, including vaccine antigens.

However, technical challenges remain that must be overcome before
20 plant-based production of complex therapeutic proteins for human and animal use gains widespread acceptance in the commercial arena. Optimization of protein production levels, an important requirement to any heterologous expression system, is one of these challenges. At present, direct expression of recombinant proteins in transgenic plants does not always satisfy the requirement for high levels of protein
25 expression. One alternative approach to expressing high levels of foreign proteins in plants is to use plant pathogens, such as plant viruses. Viral vector technology has the following advantages: (1) the flexibility to integrate and evaluate gene constructs encoding a new, improved or modified product in a few weeks' time; (2) the ability to make a range of products, from peptides to complex glycosylated proteins; and (3) the

demonstrated ability to achieve substantially lower cost of goods and services over alternative protein production systems.

Although plant virus-based vectors have great potential for producing foreign proteins, there is a need for improvement of several characteristics. Insertion of foreign sequences may result in failure or reduction of infectivity of virus due to interference with movement, assembly, or replication. Some of these difficulties may be circumvented by reducing the selective pressure of the host plant on virus movement and replication. The present invention accomplishes this reduction of selective pressure on the host plant by complementation of certain functions using transgenic plants and/or recombinant viruses.

SUMMARY OF THE INVENTION

The present invention relates generally to methods and compositions for producing foreign sequences and polypeptides in host plants using viruses. The present invention facilitates the production of desired proteins and polypeptides using transcomplementation systems involving recombinant plant viral vectors and/or transgenic plants expressing viral genes of a selected virus. Thus, the present invention is directed to a method of producing proteins and polypeptides by utilizing modified plant viruses including chimeric viruses that infect transgenic or nontransgenic plants, thereby leading to expression of the desired proteins or polypeptides throughout the plant. The terms foreign polypeptide (or protein) encoding nucleic acid sequences and heterologous nucleic acid sequences are used interchangeably herein.)

Various methods for producing foreign polypeptides in a host plant using a virus can be contemplated, as would be clear to one of ordinary skill in the art once having been apprised of the teachings.

Accordingly, the present invention provides a method for producing foreign polypeptides in a transgenic host plant through functional transcomplementation of a

virus, the method comprising: (a) constructing a recombinant viral vector for systemic infection which comprises a recombinant genomic component of the virus comprising a movement protein encoding nucleic acid sequence and a coat protein nucleic acid sequence, and one or more heterologous nucleic acid sequences cloned into the
5 recombinant genomic component such that the expression of the recombinant genomic component also results in the expression of at least one of said heterologous nucleic acid sequences; (b) providing said plant that is transgenic for expressing replicase genes of a virus, wherein said plant expressing said replicase genes complements the virus replicase function; (c) infecting said plant at one or more locations with the
10 recombinant viral vector such that the infection of said plant with the recombinant viral vector at one location results in systemic infection in the host plant; and (d) producing said foreign polypeptides in said plant infected with the recombinant viral vector by growing said plant. Alternatively, the same method can be used to produce polypeptides in a non-transgenic plant.

15 The selected genomic component can be of mono-, bi-, tri-partite genomic virus. The genomic component has a movement protein encoding gene and/or a coat protein encoding gene. At least one said foreign polypeptide-encoding nucleic acid sequence (heterologous nucleic acid sequence) which encodes a foreign polypeptide of interest is cloned into the full-length genomic component to create N-terminal fusion
20 with the coat protein. At least one foreign polypeptide-encoding nucleic acid sequence in the recombinant viral vector is an in vitro transcription promoter sequence that is placed upstream of the remaining recombinant genomic component.

The present invention also provides method for producing foreign polypeptides in a transgenic host plant through functional transcomplementation of a virus, the
25 method comprising: (a) constructing a recombinant viral vector for systemic infection which comprises a recombinant genomic component of the virus comprising a movement protein encoding nucleic acid sequence and a functional mutant coat protein nucleic acid sequence encoding an amino acid sequence having N-terminal

deletions of up to 12 amino acids, and one or more heterologous nucleic acid sequences cloned into the recombinant genomic component wherein one of said heterologous nucleic acid sequences is fused to the N-terminus of the functional mutant coat protein nucleic acid sequence such that the expression of the recombinant genomic component also results in the expression of fused heterologous nucleic acid sequence; (b) providing said plant that is transgenic for expressing replicase genes of a virus, wherein said plant expressing said replicase genes complements the virus replicase function; (c) infecting said plant at one or more locations with the recombinant viral vector such that the infection of said plant with the recombinant viral vector at one location results in systemic infection in the host plant; and (d) producing said foreign polypeptides in said plant infected with the recombinant viral vector by growing said plant.

The amino acid sequence of the functional mutant coat protein can have 1-12 amino acids deleted from the N-terminus and the foreign polypeptide-encoding nucleic acid sequence that is fused to the N-terminus of the functional mutant coat protein. The heterologous nucleic acid sequence can encode a vaccine antigen that is selected from the group consisting of hepatitis B surface antigen, enterotoxin, rabies virus glycoprotein, rabies virus nucleoprotein, Norwalk virus capsid protein, gastrointestinal cancer antigen, G protein of Respiratory Syncytial Virus, Sandostatin or colorectal cancer antigen.

The present invention yet also provides a method for producing foreign polypeptides in a transgenic host plant through functional transcomplementation of a virus, the method comprising: (a) constructing a first recombinant viral vector for infection which comprises a recombinant genomic component of the virus comprising the native movement protein encoding nucleic acid sequence and a heterologous nucleic acid sequence in place of the native coat protein encoding nucleic acid sequence such that the expression of the recombinant genomic component also results in the expression of the heterologous nucleic acid sequence; (b) constructing a second

recombinant viral vector for infection which comprises a recombinant genomic component of the virus comprising the native coat protein encoding nucleic acid sequence and a heterologous nucleic acid sequence in place of the native movement protein encoding nucleic acid sequence such that the expression of the recombinant genomic component also results in the expression of the heterologous nucleic acid sequence; (c) providing said plant that is transgenic for expressing replicase genes of a virus, wherein the transgenic plant expressing said replicase genes complements the virus replicase function; (d) infecting said plant at one or more locations with a mixture of the first recombinant viral vector and a second recombinant viral vector such that the infection of said plant with the first, second and third recombinant viral vectors at one location results in systemic infection in said plant, wherein the first recombinant viral vector expressing the native movement protein complements the cell-to-cell movement function of the virus and the second recombinant vector expressing the native coat protein complements the long distance transport function of the virus; and (e) producing said foreign polypeptides in the host plant infected with the recombinant viral vector by growing the host plant.

The present invention further provides a method for producing foreign polypeptides in a host plant through functional transcomplementation of a chimeric virus, the method comprising: (a) constructing a recombinant viral vector for systemic infection which comprises a recombinant genomic component of a first class of virus comprising a movement protein encoding nucleic acid sequence of the first class of virus, a full-length coat protein nucleic acid sequence of a second class of virus in place of the native coat protein nucleic acid sequence of the first class of virus and one or more heterologous nucleic acid sequences cloned into the recombinant genomic component such that the expression of the recombinant genomic component also results in the expression of at least one of said heterologous nucleic acid sequences; (b) infecting the host plant at one or more locations with the recombinant viral vector such that the infection of the host plant with the recombinant viral vector at one

location results in systemic infection in the host plant, wherein the recombinant viral vector expressing the native movement protein of the class of virus complements the cell-to-cell movement function of the chimeric virus and that expressing the full-length coat protein nucleic acid sequence of the different class of virus complements the long distance transport function of the chimeric virus; and (c) producing said foreign polypeptides in the host plant infected with the recombinant viral vector by growing the host plant.

The present invention also provides a method for producing foreign polypeptides in a host plant through functional transcomplementation of a chimeric virus, the method comprising: (a) constructing a first recombinant viral vector for systemic infection which comprises a recombinant genomic component of a first class of virus comprising a movement protein encoding nucleic acid sequence of the first class of virus, a non-functional coat protein nucleic acid sequence of the first class of virus, a full-length coat protein nucleic acid sequence of a second class of virus inserted into the non-functional coat protein nucleic acid sequence of the first class of virus; (b) constructing a second recombinant viral vector for infection which comprises a recombinant genomic component of a class of virus comprising a movement protein encoding nucleic acid sequence of the class of virus, a non-functional coat protein nucleic acid sequence of the class of virus, and a first heterologous nucleic acid sequences cloned into the recombinant genomic component such that the expression of the recombinant genomic component also results in the expression of the first heterologous nucleic acid sequence; (c) constructing a third recombinant viral vector for systemic infection which comprises a recombinant genomic component of the first class of virus comprising a movement protein encoding nucleic acid sequence of the first class of virus, a non-functional coat protein nucleic acid sequence of the first class of virus, and a second heterologous nucleic acid sequence together with the 5' and 3' non-coding sequences of the full-length coat protein nucleic acid sequence of the second class of virus inserted into the non-

functional coat protein nucleic acid sequence of the first class of virus such that the expression of the recombinant genomic component also results in the expression of the heterologous nucleic acid sequence; (d) infecting the host plant at one or more locations with the first, second and third recombinant viral vectors such that the infection of the host plant with said recombinant viral vectors at one location results in systemic infection in the host plant, wherein the first, second and third recombinant viral vectors expressing the native movement protein of the first class of virus complements the cell-to-cell movement function of the chimeric virus and the first recombinant viral vector expressing the full-length coat protein nucleic acid sequence of the second class of virus complements the long distance transport function of the chimeric virus; and (e) producing said foreign polypeptides in the host plant infected with the recombinant viral vector by growing the host plant.

The present invention further provides a method for eliciting an immunological response in a mammal comprising the step of: administering to the mammal an amount of a polypeptide containing plant or plant tissue thereof produced according to the methods described herein to induce an immunological response to said polypeptide in said individual.

The present invention also contemplates various compositions. Accordingly, one embodiment of the invention is a composition comprising a recombinant chimeric viral vector capable of systemic infection for producing foreign polypeptides in a host plant which comprises a recombinant genomic component of a first class of virus comprising a movement protein encoding nucleic acid sequence of the first class of virus, a coat protein nucleic acid sequence of a second class of virus in place of the native coat protein nucleic acid sequence of the first class of virus and one or more heterologous nucleic acid sequences cloned into the recombinant genomic component of the first class of virus such that the expression of the recombinant genomic component also results in the expression of at least one said heterologous nucleic acid sequence.

Another embodiment of the invention is a composition comprising a mixture of recombinant viral vectors capable of systemic infection for producing foreign polypeptides in a host plant, the mixture comprising: (a) a first recombinant viral vector which comprises a recombinant genomic component of the virus comprising a movement protein encoding nucleic acid sequence and a heterologous nucleic acid sequence in place of the native coat protein encoding nucleic acid sequence such that the expression of the recombinant genomic component also results in the expression of the heterologous nucleic acid sequence; and (b) a second recombinant viral vector which comprises a recombinant genomic component of the virus comprising the native coat protein encoding nucleic acid sequence and a heterologous nucleic acid sequence in place of the native movement protein encoding nucleic acid sequence such that the expression of the recombinant genomic component also results in the expression of the heterologous nucleic acid sequence.

An additional of the embodiment of the invention is a composition comprising a plant tissue having foreign polypeptides produced according to the methods described herein.

Another embodiment of the present invention is a recombinant in vitro transcript of an alfalfa mosaic virus for producing foreign polypeptides in a host plant, the recombinant in vitro transcript comprising: (a) a movement protein encoding nucleic acid sequence; (b) a coat protein encoding nucleic acid sequence; and (c) a heterologous protein encoding nucleic acid sequence inserted into the recombinant genomic component such that the expression of the movement protein encoding nucleic acid sequence and the coat protein encoding nucleic acid sequence also results in the expression of the heterologous nucleic acid sequence, wherein the host plant is a transgenic plant that is transgenic for expressing replicase genes of said virus, wherein the transgenic plant expressing replicase genes transcomplements the replicase function of said virus.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Schematic representation of cloning of foreign peptides as a translational fusions of AIMV CP in full-length RNA3 of AIMV.

Figure 2. Schematic representation of cloning of foreign peptides as a translational fusions with mutant AIMV CP (CPATG3) in full-length RNA3 of AIMV.

Figure 3. Schematic representation showing the construction of RNA3a vectors. RNA3 is wild-type genomic RNA of AIMV.

Figure 4. Schematic representation showing the construction of RNA3b vectors. RNA3 is wild-type genomic RNA of AIMV.

Figure 5. Western analysis and Coomassie staining of NF1/RSV.

Figure 6. Western analysis and Coomassie staining of NF2/RSV.

Figure 7. Western analysis and Coomassie staining of NF2/Sand.

Figure 8. Western analysis of NF1/TVE and NF2/TVE accumulation and assembly in infected plants.

Figure 9. Schematic representation of the genome of Av (derivative of TMV) and construction of Av/A4 (A), Av/GFP (B), and Av/A4GFP (C): the 126 kD and 183 kD proteins are required for the TMV replication, 30 kD protein is the viral movement protein, and CP is viral coat protein.

Figure 10. ELISA analysis of CO17-1A self-assembly in virus-infected plants.

Figure 11. Western analysis of P3/17-1ACH expression in plants.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed, among other things, to the methods for a novel means of production of recombinant foreign polypeptides and RNA sequences in plants using viruses. In particular, the embodiments of the methods disclosed herein use recombinant viral vectors which are capable of infecting a suitable host

plant and systemically transcribing or expressing foreign sequences or polypeptides in the host plant.

The present invention is also directed to compositions and recombinant in vitro transcripts which are capable of systemically transcribing or expressing foreign
5 sequences or polypeptides in a suitable host plant. Accordingly, in accordance with the subject invention, methods and compositions are provided for a novel means of production of foreign polypeptides and RNA sequences that can be easily separated from host cell components.

The practice of the present invention employs, unless otherwise indicated,
10 conventional techniques of molecular biology, microbiology, virology, recombinant DNA technology, and immunology particularly, which are within the skill of the art. Such techniques are explained fully in the literature. All publications and references, including but not limited to patents, cited in this specification, are herein incorporated by reference in their entirety as if each individual publication or reference were
15 specifically and individually indicated to be incorporated by reference herein as being fully set forth.

In accordance with one aspect of the subject invention, methods are provided for a novel means of production of foreign polypeptides in plants using recombinant viral vectors capable of systemically expressing foreign polypeptides upon infection.
20 Infection as used herein is the ability of a the recombinant viral vector(s) to transfer nucleic acid to a host or introduce viral nucleic acid into a host, wherein the viral nucleic acid is replicated, both viral proteins and foreign sequences are synthesized, and new viral particles assembled having foreign sequences or proteins. The foreign polypeptide of interest in the present invention is not naturally found in the host plant.

25

As a general rule, the methods of the invention require constructing one or more recombinant viral vectors to carry one or more heterologous nucleic acid sequences of interest for systemic infection in the host plants. Systemic infection or

the ability to spread systemically of a virus is the ability of the virus to spread from cell to cell and to replicate and express in most of the cells of the plant. Thus the ability to introduce nucleic acid into one part of a plant and have it spread to the rest of the plant would overcome the problems of growing plants from transgenic cultures.

- 5 The methods of the invention can also require that after infecting a host plant with one or more recombinant viral vectors, heterologous nucleic acid sequences of interest are expressed systemically in host plants by complementation of certain functions provided by the host plants that are transgenic for certain viral genes and/or the recombinant viral vectors. The complementation functions provided by the host plants
- 10 and/or the recombinant viral vectors include virus replication, assembly and movement (cell-to-cell or long distance movement). The systemic spread of foreign sequences or polypeptides through complementation functions provided by the host plants and/or the recombinant viral vectors is an essential feature of the invention. In some aspects of the invention, the recombinant viral vectors are designed such that the
- 15 heterologous nucleic acid sequences are expressed systemically through transcomplementation.

- Thus, the subject method includes the steps of constructing a recombinant viral vector having two or more heterologous nucleic acid sequences, infecting the host plant with the recombinant viral vector and producing foreign polypeptide of interest
- 20 in the host plant by allowing the host plant to grow for some time. The process can also include isolating the desired product, if necessary. The growth of the infected host is in accordance with conventional techniques as is the isolation of the desired product. Purification of the recombinant protein, if required, is greatly simplified. The recombinant DNA or RNA encoding the polypeptide of interest can be part or all
- 25 of a naturally occurring gene from any source, it may be a synthetic DNA or RNA sequence or it may be a combination of naturally occurring and synthetic sequences.

Thus, the first step in achieving any of the features of the invention is to construct a recombinant viral vector by manipulating the genomic component of a

virus. Preferred virus is RNA containing plant virus. Although many plant viruses have RNA genomes, it is well known that organization of genetic information differs among groups. Thus, a virus can be a mono-, di-, tri-partite virus. "Genome" refers to the total genetic material of the virus. "RNA genome" states that as present in virions
5 (virus particles), the genome is in RNA form.

Some of the viruses which meet this requirement, and are therefore suitable, include Alfalfa Mosaic Virus (AIMV), ilarviruses, cucumoviruses such as Cucumber Green Mottle Mosaic virus (CGMMV), closteroviruses or tobamaviruses (tobacco mosaic virus group) such as Tobacco Mosaic virus (TMV), Tobacco Etch Virus
10 (TEV), Cowpea Mosaic virus (CMV), and viruses from the brome mosaic virus group such as Brome Mosaic virus (BMV), broad bean mottle virus and cowpea chlorotic mottle virus. Additional suitable viruses include Rice Necrosis virus (RNV), and geminiviruses such as tomato golden mosaic virus (TGMV), Cassava latent virus (CLV) and maize streak virus (MSV). Each of these groups of suitable viruses are
15 well characterized and are well known to the skilled artisans in the field.

It should be noted that chimeric genes and vectors and recombinant plant viral nucleic acids according to this invention are constructed using techniques well known in the art. Briefly, manipulations, such as restriction, filling in overhangs to provide blunt ends, ligation of linkers, or the like, complementary ends of the fragments can be
20 provided for joining and ligation. In carrying out the various steps, cloning is employed, so as to make the desired virus genomic component and heterologous nucleic acid combinations, to amplify the amount of DNA and to, allow for analyzing the DNA to ensure that the operations have occurred in proper manner. A wide variety of cloning vectors are available, where the cloning vector includes a replication system
25 functional in *E. coli* and a marker which allows for selection of the transformed cells. Illustrative vectors include pBR332, pUC series, M13mp series, pACYC184, etc for manipulation of the primary DNA constructs. See Life Technologies Catalogue (1999); Amersham Pharmacia Biotech Catalogue (1999). Thus, the sequence may be

inserted into the vector at an appropriate restriction site(s), the resulting plasmid used to transform the *E. coli* host, the *E. coli* grown in an appropriate nutrient medium and the cells harvested and lysed and the plasmid recovered. Analysis may involve sequence analysis, restriction analysis, electrophoresis, or the like. After each
5 manipulation the DNA sequence to be used in the final construct may be restricted and joined to the next sequence, where each of the partial constructs may be cloned in the same or different plasmids. Suitable techniques have been described in standard references and well known to one skilled in the art. DNA manipulations and enzyme treatments are carried out in accordance with manufacturers' recommended
10 procedures.

Cloning of heterologous nucleic acid sequences into the selected recombinant genomic component of the virus can take place in various ways including terminal fusions (N-terminal, C-terminal) and/or internal fusions. Construction of fusion protein requires the identification of a suitable restriction site close to the translational
15 start codon of a gene of the viral vector, a coat protein gene for example. A suitable restriction site can be created without any alteration in coding sequence by the introduction of base changes in the start codon. As an illustration of such a modification is the A1MV coat protein shown in Figure 1A. The A1MV coat protein here is modified in such a way that replacement of AU in AUG by TC yields an *XhoI*
20 site. Alternatively, other restriction sites may be used or introduced to obtain cassette vectors that provide a convenient means to introduce heterologous nucleic and sequences encoding foreign polypeptide. The coding sequence for the foreign polypeptide can require preparation which will allow its ligation directly into the created site in the viral vector. For example, introduction of a foreign polypeptide
25 encoding sequence into the *XhoI* site introduced into the A1MV coat protein described above can require the generation of compatible ends for ligation. This can typically require a single or two-base modification of site-directed mutagenesis to generate *XhoI* around C-terminus of the foreign peptide. The preferred method would be to use

primers as linkers to produce the foreign polypeptide encoding sequence flanked by appropriate restriction sites. Orientation is checked by the use of restriction sites in the coding sequence.

5 The resultant construct from these N-terminal fusions would contain AIMV coat protein promoter sequence, an in-frame fusion in the first few condons of the AIMV coat protein gene of a desired foreign polypeptide-encoding sequence with its own ATG as start signal and the remainder of the AIMV protein gene sequence and terminator. Thus, protein synthesis can occur in the usual way, from the starting
10 codon for methionine on the foreign gene to the stop codon on the viral gene (e.g., coat protein) to produce the fusion protein. In all fusions, the regulation sites on the viral genome can remain functional. Foreign polypeptide or protein-encoding nucleic acid sequence as used herein refers to the sequences that encode foreign polypeptide or protein of interest such as for example, vaccine antigen, antibodies etc.

15 Internal fusions involve placing of the foreign polypeptide encoding sequences or the coat protein encoding sequences of a different class of virus internally to the coding sequence of the virus, e.g., coat-protein encoding sequence. Thus, various strategies are dependent on the particular use of the nucleic acid sequence of the foreign polypeptide and would be apparent to those skilled in the art.

20 In some embodiments, the nucleic acid sequences encoding the foreign polypeptides or proteins (cargo peptide) are further engineered for generating recombinant polypeptides or proteins with inherent cell membrane-translocating activity in animals. Typically, in designing such a recombinant polypeptide or protein, a region of a signal peptide (used as a carrier for import into animal cells) can be placed at either the N-terminus or the C-terminus of the polypeptide of interest i.e.,
25 cargo peptide. Using this strategy it is possible to bestow membrane-translocating ability on a wide variety of proteins ranging from few amino acids to large molecular masses (over 40 kDa) to allow efficient import into living cells. When administered in the animals. It is well known in the art that the hydrophobic region (h region) of a

signal peptide sequence can be used as a carrier to deliver peptides (cargo) into living cells without destroying their activity. See Lin et al., 1995, J. Biol. Chem., 270: 14255-14258. Thus foreign proteins (cargo peptides), particularly vaccine antigens and antibodies can be made cell-membrane permeable simply after its attachment of
5 fusion to a short membrane-translocating peptide sequence.

These plant produced foreign polypeptides of interest can be vaccine antigens which can be administered parenterally and/or orally. These vaccine antigens can also be engineered to fuse with proteins such as protective antigen of anthrax bacteria, heat shock protein which can facilitate the transport of administered antigen to cell cytosol.
10 It should be noted that the vaccine antigens can be co-expressed and co-administered together with immunoenhancers such as cytokines and hormones.

The viral coat protein gene need not encode a full-length protein; any encoded coat protein that acts as a carrier molecule of the fused protein and retains the encapsidation function is sufficient. Numerous methods are known to one skilled in
15 the art to delete sequence from or mutate nucleic acid sequences that encode a protein and to confirm the function of the proteins encoded by these deleted or mutated sequences. Accordingly, the invention also relates to a mutated or deleted versions of a coat protein nucleic acid sequence (analogs or mutant coat proteins) of viral genomic component that encodes a protein that retains a known function. These analogs can
20 have N-terminal, C-terminal or internal deletions, so long as function is retained. The inventors of the present invention discovered that the maximum number of amino acids that can be deleted from the N-terminus of the AIMV coat protein without altering its function is 14 amino acids. In some instances the coat protein carrying such deletions can perform significantly better than the full-length protein. Thus, in
25 particular preferred embodiments of the invention the coat protein of a virus can have the first 10 to 12, 5 to 10, 1 to 5, 1 to 4, 3, 2 or up to 12 or up to 14 amino acid residues deleted from the N-terminus of the coat protein. In some other embodiments the first 10 to 12, 5 to 10, 1 to 5, 1 to 4, 3, 2, 1 or up to 12 or up to 14 amino acid

residues of the N-terminus of the coat protein are modified or substituted in any combination. Especially preferred among these are silent substitutions and/or modifications that either enhance or do not alter the properties and activities of the resulting coat protein when present in the viral vectors used to infect plants.

- 5 Deletional and modification approaches can also be applied to the other nucleic acid sequences of a virus such as the movement protein encoding sequences.

The transcription termination region which is employed will be primarily one of convenience, since in many cases termination regions appear to be relatively interchangeable. The transcription termination region is a sequence that controls
10 formation of the 3' end of the transcript. For example, polyadenylation sequences and self-cleaving ribozymes. The transcription termination region may be native to the transcriptional initiation region, may be native to the heterologous nucleic acid sequence encoding the polypeptide of interest, or may be derived from another source. Termination signals for expression in other organisms are well known in the
15 literature. Sequences for accurate splicing of the transcript may also be included. Examples are introns and transposons.

Recombinant viral vectors used herein can be *in vitro* transcripts. After assembly of a recombinant genomic component and heterologous nucleic acid sequence(s) encoding polypeptide(s) combination, this combination can be placed
20 behind a (downstream of) heterologous promoter (a heterologous nucleic acid sequence) that can drive *in vitro* transcription of the downstream sequences to produce *in vitro* transcripts. Examples of efficient heterologous promoters for *in vitro* transcription include a bacteriophage promoter such as the T7 phage promoter or SPG promoter. After such a viral vector/heterologous nucleic acid sequence(s) encoding
25 polypeptide(s)/*in vitro* transcription vector combination is assembled, *in vitro* transcripts for infection can be produced by *in vitro* transcription and mixed with any other viral RNA *in vitro* transcripts necessary for maintenance of the viral vector in a plant cell. RNA production from the vector can be conducted, for instance, with the

method described in Ausubel et al., *SHORT PROTOCOLS IN MOLECULAR BIOLOGY*, John Wiley & Sons, New York, 1992.

The *in vitro* transcripts for infection can be applied to recipient cell(s) of a plant by any of the techniques known to those skilled in the art. Suitable techniques include, but are not limited to, hand inoculations such as abrasive inoculations (leaf abrasion, abrasion in a buffer solution), mechanized spray inoculations, vacuum infiltration, particle bombardment and/or electroporation. It should be realized that the use of a mixture viral vectors can depend on the type of plant host and/or the class of virus vector used for infections. Thus in one strategy, where a Δ IMV virus replicase expressing transgenic plant host is used, then a mixture of recombinant viral vectors each having AIMV RNA3 or RNA4 genomic components is used. In another strategy, for example, where a non-transgenic plant host and AIMV virus as a viral vector are used, then a mixture of the recombinant viral vectors each having AIMV RNA1, RNA2, RNA3 or RNA4 is used.

Suitable buffer solutions in which the recombinant vectors are suspended to prepare inoculum for inoculation are well known in the art. For example, leaves of plants can be inoculated with *in vitro* transcription products of recombinant viral vectors as described (Yusibov et. al., 1997) after adding 1 vol (v/v) of FES buffer [sodium-pyrophosphate 1% (w/v), malacoid 1% (w/v), celite 1% (w/v), glycine 0.5 M, K_2HPO_4 0.3 M, pH 8.5, with phosphoric acid]. The mixture *in vitro* transcription products and FES buffer can be applied to leaves after abrading the leaf surface with carborundum (320 grit; Fisher, Pittsburgh, PA). Inoculation can be affected by gentle rubbing to spread the inoculum and further abrade the leaf surface.

In preferred aspects of the invention, the initial plant inoculation can be carried out with *in vitro* transcripts. Once the recombinant virus particles are harvested from the host plant, these virus particles can be used as stock for further inoculations without having to use *in vitro* transcripts.

For embodiments of the invention wherein the foreign polypeptides are produced in plants using recombinant viral vectors, all or most of the cis-active sequences of the wild type virus which encode components necessary for production of viral particles are retained. In these embodiments heterologous nucleic acid sequences that encode foreign polypeptides are cloned into the recombinant genomic component. For example, a heterologous nucleic acid sequence that encodes a foreign polypeptide can be inserted into the recombinant genomic component of a virus having both movement protein and coat protein sequences (a situation where cis-active sequences of the wild type virus are retained).

In some instances, however, cis-active sequences which encode components necessary for production of viral particles are optionally deleted from or are rendered inactive in the recombinant viral vectors. In these instances, the missing components are supplied by complementation. For example, the missing components can be supplied by complementation in cis or in trans from a second recombinant viral vector. It is well known in the art that the term "in cis" indicates that two sequences are positioned on the same strand of RNA or DNA. The term "in trans" indicates that two sequences are positioned on different strands of RNA or DNA.

Thus, in one preferred embodiment the plant is infected with more than one recombinant viral vector (co-infection) each of which has a complementary role in the production of a viral particle. For example, the coat protein gene in a first recombinant viral vector is replaced by a heterologous nucleic acid sequence encoding a foreign polypeptide. In a second recombinant viral vector the movement protein gene is replaced by a different or same heterologous nucleic acid sequence encoding a foreign polypeptide as in the first recombinant viral vector. The first and second recombinant viral vectors can be mixed for co-infection as complementary vectors for transcomplementation. See also, the Examples. In another preferred embodiment, other functions for viral particle formation or propagation not supplied by transcomplementation described in the above embodiment are present in host plants

transgenic for viral genes. For example transgenic plants expressing viral replicase genes, i.e., ReP plants, can be used to complement the virus replicase function.

An unexpected aspect of the present invention is the discovery that the coat protein gene of a first class of virus (eg., TMV) ciscomplements the long distance
5 movement and encapsidation functions of a second class of virus (eg., AIMV) (See Example 5). Thus the complementation in the present invention can be applied rather broadly across various strains and even various genera including viruses and plants. The complementation of certain functions thus achieved has the advantage in, among other things, reducing the selective pressure by the host plant thereby facilitating the
10 movement, assembly, or replication of the recombinant viral vectors and in extending plant host range of the recombinant viruses.

A variety of techniques are available for the genetic transformation of plants and plant tissues (i.e., the stable integration of foreign DNA into plants) and are well-known to those skilled in the art. These include transformation by *Agrobacterium*
15 species and transformation by direct gene transfer. For example, the chimeric DNA constructs may be introduced into host cells obtained from dicotyledonous plants, such as tobacco and brassicas using standard *Agrobacterium* vectors by a transformation protocol such as that described by Moloney et al., 1989, Plant Cell Rep., 8:238-242 of Hinchee et al., 1988, Bio/Technol., 6:915-922; or other techniques known to those
20 skilled in the art. For example, the use of T-DNA for transformation of plant cells has received extensive study and is amply described in Knauf, et al., (1983), Genetic Analysis of Host Range Expression by *Agrobacterium*, p. 245, In: Molecular Genetics of the Bacteria-Plant Interaction, Puhler, A. ed., Springer-Verlag, NY; Hoekema et al., (1985), Chapter V, In: The Binary Plant Vector System Offset-drukkerij Kanters B.
25 V., Alblasterdam; and An et al., (1985), EMBO J., 4:277-284. Briefly, explants can be co-cultivated with *A. tumefaciens* or *A. rhizogenes* to allow for transfer of the transcription construct to the plant cells. Following transformation using *Agrobacterium*, the plant cells are dispersed in an appropriate medium for selection,

subsequently callus, shoots and eventually plantlets are recovered. The *Agrobacterium* host will harbour a plasmid comprising the vir genes necessary for transfer of the T-DNA to the plant cells. See also, Dodds, J. ed., Plant Genetic Engineering, Cambridge University Press, Cambridge (1985).

5 The use of non-*Agrobacterium* techniques permits the use of the constructs described herein to obtain transformation and expression in a wide variety of monocotyledonous and dicotyledonous plants and other organisms. These techniques are especially useful for species that are intractable in an *Agrobacterium* transformation system. Other techniques for gene transfer include biolistics (Sanford, 10 1988, Trends in Biotech., 6:299-302), electroporation (Fromm et al., 1985, Proc. Natl. Acad. Sci. U.S.A., 82:5824-5828; Riggs and Bates, 1986, Proc. Natl. Acad. Sci. U.S.A. 82:5602-5606 or PEG-mediated DNA uptake (Potrykus et al., 1985, Mol. Gen. Genet., 199:169-177).

15 The foreign polypeptides of interest to be produced using viruses by any of the specific methods described herein, can be any peptide or protein. The heterologous nucleic acid sequence encoding the polypeptide of interest can be naturally derived, synthetic, or a combination thereof.

20 The invention is not limited by the source or the use of the recombinant polypeptide. Of particular interest are those proteins or peptides that have a biomedical, therapeutic and/or diagnostic value. These proteins or polypeptides include vaccine antigens, such as viral coat proteins or G proteins or microbial cell wall or toxin proteins, cancer antigens or various other antigenic peptides, antibodies, specifically a single-chain antibody having a translational fusion of the VH or VL chains of an immunoglobulin, peptides of direct therapeutic value such as interleukin- 25 1, the anticoagulant hirudin and blood clotting factors. Vaccine antigens derived from pathogenic parasites such as *Entamoeba* and the like can also be used. Other biomedical agents such as human growth hormone or bovine somatotropin can also be produced.

In particular, the vaccine agents from the following pathogens can be particularly mentioned; *S. typhi* (the cause of human typhoid), *S. typhimurium* (the case of salmonellosis), *S. enteritis* (a cause of food poisoning in humans), *S. cholerae* (the cause of salmonellosis in animals), *Bordetella pertussis* (the case of whooping
5 cough), *Haemophilus influenzae* (a cause of meningitis), *Neisseria gonorrhoeae* (the cause of gonorrhoea) and *Haemophilus*. The vaccine agents from pathogenic parasites such as *Entamoeba* are also included.

In accordance with the present invention, the host plants included within the scope of the present invention are all species of higher and lower plants of the Plant
10 Kingdom. Mature plants, seedlings, and seeds are included in the scope of the invention. A mature plant includes a plant at any stage in development beyond the seedling. A seedling is a very young, immature plant in the early stages of development. Specifically, plants that can be used as hosts to produce foreign sequences and polypeptides include and are not limited to Angiosperms, Bryophytes
15 such as Hepaticae (liverworts) and Musci (mosses); Pteridophytes such as ferns, horsetails, and lycopods; Gymnosperms such as conifers, cycads, Ginkgo, and Gnetales; and Algae including Chlorophyceae, Phaeophyceae, Rhodophyceae, Myxophyceae, Xanthophyceae, and Euglenophyceae.

Plants for the production of desired sequences can be grown either in vivo
20 and/or in vitro depending on the type of the selected plant and the geographic location.

It is important that the selected plant is plant amenable to cultivation under the appropriate field conditions and/or in vitro conditions. The conditions for the growth of the plants are described in various basic books on botany. Agronomy, Taxonomy and Plant Tissue Culture, and are known to a skilled artisan in these fields.

25 Among angiosperms, the use of crop and/or crop-related members of the families identified in the paragraph below are particularly contemplated. The plant members used in the present methods also include interspecific and/or intergeneric hybrids, mutagenized and/or genetically engineered plants. The term "crop member"

refers specifically to species which are commercially grown as sources for vegetables, grains, forage, fodder, condiments and oilseeds. Crop-related" members are those plants which have potential value as a crop and as donors of agronomically useful genes to crop members. Thus, crop-related members are able to exchange genetic material with crop members, thus permitting breeders and biotechnologists to perform interspecific (i.e., from one species to another) and intergeneric (i.e., from one genus to another) gene transfer. Those having ordinary skill in the art will understand that methods of exchanging genetic material between plants and testing effects of interspecific and intergeneric gene transfer are well characterized. See, for example Goodman et al., Science, 236: 48-54, 1987, incorporated herein by reference.

These families include and not limited to Leguminosae (Fabaceae) including pea, alfalfa, and soybean; Gramineae (Poaceae) including rice, corn, wheat; Solanaceae particularly of the genus *Lycopersicon*, particularly the species *esculentum* (tomato), the genus *Solanum*, particularly the species *tuberosum* (potato) and *melongena* (eggplant), the genus *Capsicum*, particularly the species *annuum* (pepper), tobacco, and the like; Umbelliferae, particularly of the genera *Daucus*, particularly the species *carota* (carrot) and *Apium*, particularly the species *graveolens dulce*, (celery) and the like; Rutaceae, particularly of the genera *Citrus* (oranges) and the like; Compositae, particularly the genus *Lactuca*, and the species *sativa* (lettuce), and the like and the Family Cruciferae, particularly of the genera *Brassica* and *Sinapis*. Examples of "vegetative" crop members of the family Brassicaceae include, but are not limited to, digenomic tetraploids such as *Brassica juncea* (L.) Czern. (mustard), *B. carinata* Braun (ethopian mustard), and monogenomic diploids such as *B. oleracea* (L.) (cole crops), *B. nigra* (L.) Koch (black mustard), *B. campestris* (L.) (turnip rape) and *Raphanus sativus* (L.) (radish). Examples of "oil-seed" crop members of the family Brassicaceae include, but are not limited to, *B. napus* (L.) (rapeseed), *B. campestris* (L.), *B. juncea* (L.) Czern. and *B. tournifortii* and *Sinapis alba* (L.) (white mustard).

In sum, the plant members used in the present invention are plants that: (a) can be grown to high biomass in a short time either in vivo or in vitro; (b) are adaptable for growth in various agroclimatic conditions; (c) are adaptable to modified, non-conventional agricultural practices, described herein, for monoculture; (d) are
5 amenable to genetic manipulation by mutagenesis and/or gene transfer; and (e) can produce several crops per year. Additionally the plant members are natural hosts for a selected virus. Alternatively, the selected virus can be made compatible with a plant so as to function as a host.

Depending on the type of host plant (lower plants to higher plants in the plant
10 kingdom) used, infected or systemically infected host plant or tissue thereof can be harvested 10 days after inoculation, preferably 14 days after inoculation and more preferably 16 days after inoculation. Samples for the analysis (detection and quantification) of recombinant viruses and desired sequences can be taken from crude extracts of infected plant and from purified recombinant virus. Recombinant viruses can
15 be purified from infected tissue can be easily accomplished using standard virus purification procedures known in the art.

Polypeptides and polynucleotides of interest can be recovered and purified from recombinant viruses by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography,
20 phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and lectin chromatography. High performance liquid chromatography can be employed for purification. Well known techniques for refolding protein can be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

25 Purification techniques other than the affinity procedures outlined above can be used to purify, or supplement the purification of, a protein of the invention. Such methods can include without limitation, preparative electrophoresis, FPLC (Pharmacia, Uppsala, Sweden), HPLC (e.g., using gel filtration, reverse-phase or

mildly hydrophobic columns), gel filtration, differential precipitation (for instance, "salting out" precipitations) and ion-exchange chromatography. The matrix used to create the affinity matrices will preferably comprise a carbohydrate matrix such as cross-linked dextran (e.g., that sold under the tradename Sepharose) or agarose (e.g., that sold by Pharmacia, Sweden as "Sephacryl"). The matrix should have pore sizes sufficient to admit both the affinity ligand that will be attached to the matrix and the multifunctional enzyme of the invention. Methods of synthesizing appropriate affinity columns are well known. See, for instance, Axén et al., *Nature*, 214:1302-1304, 1967.

The polypeptides and nucleic acids in the recombinant viruses are detected and quantified by any of a number of means well known to those of skill in the art. The infected plants can show symptoms specific to each virus. Such symptom production can be a useful detection marker. A number of laboratory techniques can also reliably be employed for the detection. These include analytic biochemical methods such as spectrophotometry, radiography, electrophoresis, capillary electrophoresis, high performance liquid chromatography (HPLC), thin layer chromatography (TLC), hyperdiffusion chromatography, and the like, and various immunological methods such as fluid or gel precipitin reactions, immunodiffusion (single or double), immunoelectrophoresis, radioimmunoassays (RIAs), enzyme-linked immunosorbent assays (ELISAs), immunofluorescent assays, and the like. The detection of nucleic acids proceeds by well known methods such as northern analysis, gel electrophoresis, PCR, radiolabeling and scintillation counting, and affinity chromatography.

One of skill will appreciate that it is often desirable to reduce non specific binding in immunoassays and during analyte purification. Where the assay involves a polypeptide, antibody, or other capture agent immobilized on a solid substrate, it is desirable to minimize the amount of non specific binding to the substrate. Means of reducing such non specific binding are well known to those of skill in the art. Typically, this involves coating the substrate with a proteinaceous composition. In

particular, protein compositions such as bovine serum albumin (BSA), nonfat powdered milk, and gelatin are widely used.

Western blot analysis can also be used to detect and quantify the presence of a transcript polypeptide or antibody or enzymatic digestion product) in the sample. The technique generally comprises separating sample products by gel electrophoresis on the basis of molecular weight, transferring the separated proteins to a suitable solid support, (such as a nitrocellulose filter, a nylon filter, or derivatized nylon filter), and incubating the sample with labeling antibodies that specifically bind to the analyte protein. The labeling antibodies specifically bind to analyte on the solid support. These antibodies are directly labeled, or alternatively are subsequently detected using labeling agents such as antibodies that specifically bind to the labeling antibody. To prevent nonspecific binding of compound (e.g., labeled antibody) to the surface of the solid support, the surface is typically blocked with a second compound (e.g., milk).

Labeling agents include e.g., monoclonal antibodies, polyclonal antibodies, proteins such as those described herein, or other polymers such as affinity matrices, carbohydrates or lipids. Detection proceeds by any known method, such as immunoblotting, western analysis, gel-mobility shift assays, fluorescent in situ hybridization analysis (FISH), tracking of bioluminescent markers, nuclear magnetic resonance, or other methods which track a molecule based upon size, charge or affinity. Thus, a label is any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical or chemical means. Useful labels in the present invention include magnetic beads (e.g. Dynabeads.TM.), fluorescent dyes (e.g., fluorescein isothiocyanate, rhodamine, and the like), radiolabels (e.g., ^3H , ^{251}I , ^{35}S , ^{14}C , or ^{32}P), and nucleic acid intercalators (e.g., ethidium bromide)

The label is coupled directly or indirectly to the desired component of the assay according to methods well known in the art. As indicated above, a wide variety of labels are used, with the choice of label depending on the sensitivity required, ease

of conjugation of the compound, stability requirements, available instrumentation, and disposal provisions.

Means of detecting labels are well known to those of skill in the art. Thus, for example, where the label is a radioactive label, means for detection include a
5 scintillation counter or photographic film as in autoradiography. Where the label is a fluorescent label, it may be detected by exciting the fluorochrome with the appropriate wavelength of light and detecting the resulting fluorescence, e.g., by fluorescence microscopy, visual inspection, via photographic film or by the use of electronic detectors and the like.

10 Animal or human hosts infected by a pathogen or vaccine antigens (or antigenic or immunogenic determinants) mount an immune response in response to the invading pathogen or the vaccine antigen. The immune system works in three fundamentally different ways which is well known in the art. See, Lodish et al., MOLECULAR CELL BIOLOGY, Scientific American Books, New York (1995);
15 Roitt et al. *IMMUNOLOGY*, Mosby International, London (1998) which are incorporated herein by reference. Accordingly, The foreign polypeptides or polynucleotides or cells expressing them produced according to the methods described herein can be used as an antigen or as an immunogen for vaccination of an animal including human to produce specific antibodies which have anti-bacterial anti-viral
20 and/or anti-cancerous action. In addition, polypeptides in which one or more of the amino acid residues are modified (i.e., derivatives of polypeptides) can also be used. Such polypeptides, for example, can be the result of substitution, addition, or rearrangement of amino acids or chemical modification thereof. All such substitutions and modifications are generally well known to those skilled in the art of peptide
25 chemistry.

This invention also contemplates the use of the foreign nucleic acids encoding the antigen as a component in a DNA vaccine as discussed further below.

Another aspect of the invention relates to a method for inducing an immunological response in an animal, particularly a human which involves administering the animal an effective amount of plant cells or tissue containing a vaccine antigen, or a purified vaccine antigen produced according to the method
5 herein, adequate to produce antibody and/ or T cell immune response to protect said animal from infection and/or disease caused by pathogens. Also provided are methods whereby such immunological response slows bacterial or viral replication or a parasitic pathogen in the animal whether that disease is already established within the animal or not.

10 Polypeptides and their derivatives include immunologically equivalent derivatives which form a particular aspect of this invention. The term 'immunologically equivalent derivative' as used herein encompasses a peptide or its equivalent which when used in a suitable formulation to raise an immunological response in an animal which response acts to interfere with the interaction between
15 pathogen and mammalian host. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity or cellular immunity such as that arising from CTL or CD4+ T cells.

The polypeptide, such as an immunologically equivalent derivative or a fusion protein thereof is used as an antigen to immunize the animal (see Example 7). The
20 fusion protein can provide stability to the polypeptide. The antigen may be associated, for example by conjugation, with an immunogenic carrier protein for example, protective antigen of anthrax bacteria and heat shock proteins which can facilitate the transport of administered antigen to cell cytosol. Alternatively, a multiple antigenic peptide comprising multiple copies of the protein or polypeptide, or an
25 immunologically equivalent polypeptide thereof may be sufficiently antigenic to improve immunogenicity so as to obviate the use of a carrier.

The invention also includes a vaccine formulation which comprises an immunogenic recombinant protein of the invention together with a suitable carrier.

Since the protein may be broken down in the stomach, it is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. In therapy or as a prophylactic, the active agent i.e., the desired vaccine antigen (polypeptide or polynucleotide) can be administered to a patient as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

As noted earlier, the plant viruses replicate and express at a high rate in plant cells, thereby leading to the rapid production of large numbers of virus particles with the attached foreign sequences following infection. Many viruses have between 10^6 to 10^7 particles per cell. Thus, there is on the order of 10-100 pg of viral coat protein per cell. Thus, according to the methods of the present invention using the recombinant viral vectors, foreign polypeptides in the order of 0.5-1.0 mg can be produced per gram of plant tissue when the foreign polypeptide is fused to the coat protein of a given virus.

The methods of producing fused coat proteins with antigenic or non-antigenic foreign polypeptides using plant viruses and methods of delivering a fused coat protein to an animal for purposes of inducing an immune response against the foreign

polypeptide has been demonstrated in WO 98/08375, the contents of which are incorporated herein by reference.

The following examples further illustrate the present invention, but of course, should not be construed as in any way limiting its scope. The examples below are carried out using standard techniques, that are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention. All animal methods of treatment or prevention described herein are preferably applied to mammals, most preferably humans.

Example 1. Construction of NF1 to express different foreign peptides.

The starting plasmid pCPΔAUG (Loesh-Fries et. al, 1985, *Virology* 146, 177-187) contains an AIMV coat protein modified so that the AUG translation initiation codon is replaced by TCG to create an XhoI (CTCGAG) site for cloning and an RNA molecule defective in translation. pSPΔAUG was used to create all NF1 constructs (Figs. 1A and B).

Shown in figure 1 is a schematic representation of cloning of foreign peptides as a translational fusions of AIMV CP in full-length RNA3 of AIMV. (A) represents cloning strategy used. Two ellipsoids represent a foreign peptide with Xho I and Sal I cloning sites. CPΔAUG is AIMV CP where the translation initiation codon (AUG) is mutated to create Xho I cloning site. T7P3Sal is 5' portion of AIMV RNA3 containing ORF for P3, 5' non coding regions of RNA3 and subgenomic RNA4. The Sal I site is created at the position 1192 to mutate the AIMV CP AUG. Additionally T7P3Sal contains T7 promoter for in vitro synthesis of infectious RNA3 transcripts. NF1 is full-length RNA3 containing foreign peptide fused to AIMV CP. (B) pNF1/g24 is RNA3 containing epitope from rabies glycoprotein. pNF1/RSV contains 24 amino acid epitope from RSV G protein. pNF1/Sand contains octerotide sandostatin and pNF1/TVE contains CP fused with 104 amino acids from colorectal

cancer GA733-2. Amino acid sequences of cloned peptides are shown under each construct. Stem-loop structure indicates the 3' non coding region of RNA3.

It should be noted that only those constructs that have the foreign polypeptide-encoding nucleic acid ligated in the right orientation is used. Right orientation of the foreign polypeptide-encoding nucleic acid is with the 5'-end thereof proximal to the promoter. Right orientation is confirmed by restriction digest analysis or by sequencing analysis.

a. **pNFI/RSV** (Fig. 1B) PCR cloning was used to create a fusion protein consisting of a full-length AIMV coat protein and the 24 amino acid peptide from the G protein of the Respiratory Syncytial Virus (RSV), shown to protect (Bastien et. al., 1997, *Virology* **234**,118-22) immunized mice against infection with RSV. DNA sequences encoding these 24 amino acids (Fig. 1B; SEQ ID NO: 1) were PCR amplified and cloned into the AIMV genome by translational fusion with the coat protein. The epitope was amplified using

5'GCCGCTCGAGCATCATGTCACCCTGCAGCATATGCAGCAACAATCCA3' (SEQ ID NO: 2) as a first strand primer and

5'CGCGTCGACTTGCAGATAGCCCAGCAGGTTGGATTGTTGCT3' (SEQ ID NO: 3) as a second strand primer. A 90-basepair (bp) PCR fragment (encoding 24 amino acids) containing the RSV epitope with introduced XhoI and SalI sites at 5'- and 3'-ends respectively, was digested with XhoI+SalI and ligated into pSPΔAUG linearized by XhoI to engineer CPRSV. The resulting plasmid is termed pSPCPRSV. Translation of the recombinant protein initiates from the AUG codon created at the 5' end of the chimeric gene, which is upstream of nucleotide sequences encoding the RSV epitope. The recombinant plasmid also contains linking 5'- (37 nucleotides upstream from the wild-type AIMV coat protein translation start codon) and 3'- (192 nucleotides following the AIMV coat protein stop codon and containing the AIMV origin of assembly) noncoding regions of the AIMV coat protein. The epitope was fused to the N-terminus of the coat protein. After sequence confirmation, the

recombinant coat protein was subcloned into full-length RNA3 of AIMV to create pNF1RSV (Fig. 1B). Using the full-length coat protein and the strategy described herein, additional constructs were engineered as follows:

b. **pNF1/g24** (Fig. 1B). PCR was performed using

- 5 5'GCGCTCGAGGGTACCATG'TCCGCCGTCTACACCCGAATTATGATGAACG
GAGGACGACTTAAGCGACCACCAGACCAGCTTG3' (SEQ ID NO: 4) as a first
strand primer and
5'CGAGGTACCTCTTCCACCACAAGGTGCTCATTTTTCGTCCGATCCGAAGT
CGTGAAGGTTTACAAGCTGGTCTGGTGGTCGCTTAAAGTCGTCC3' (SEQ ID
10 NO: 5) as a second strand primer. NF I Drg24 contains the coat protein fused with a
rabies epitope capable of protecting the immunized mice against a lethal dose of
challenge rabies virus. The next step involved the engineering of a linear epitope,
G5-24 of rabies virus glycoprotein as a chimera (Drg24) with an epitope from the
rabies virus nucleoprotein (31D). This chimera was fused with the AIMV coat protein.
15 The chimeric epitope, Drg24, was synthesized by PCR using oligonucleotides
containing 18 complementary nucleotides between the first and second, such that the
complementary nucleotide strands can anneal and initiate the PCR reaction. The 120
bp (coding for 40 amino acids; SEQ ID NO: 6) PCR product was digested with XhoI
and cloned into pSPDAUG to create pSPCPDrg24. The latter was combined by
20 ligation with the 5' part of AIMV RNA3 to obtain pNF1/g24.

c. **pNF1/Sand** (Fig. 1B). Sandostatin is an 9 amino acid peptide (SEQ ID
NO: 7) used to suppress the synthesis of human growth hormone in diseased people.

Sandostatin was fused with the coat protein of AIMV by PCR using

- 5'GCGGAATTCGTTTTTATTTTAAATTTCTTTCAATTACTTCCATCATGAGTT
25 CTTTCTGTTTCTGAAA3' (SEQ ID NO: 8) as a first strand primer and
5'GCGCTCGAGCGAGTACACGTTTTCCAGAAACAGAA3' (SEQ ID NO: 9) as a
second strand primer. The fusion product was then cloned into full-length RNA3 as
described above.

d. **pNFI/TVE** (Fig. 1B). 104 amino acid peptides from colorectal cancer antigen GA733-2 (Linnenbach et al., 1989, *Proc. Natl. Acad. Sci. USA* **86**: 27-31) were fused with mutant AIMV CP. This region has a special conformation recognized by colorectal cancer associated antibody 17-1A. The sequences encoding this region
5 (104 amino acids; SEQ ID NO: 10) of GA733-2 were PCR amplified using
5'GCGCTCGAGGGTACCATGCGACGGCGACTTTTGCCGCA3' (SEQ ID NO: 11)
as a first strand primer and
5'GTCGACCTGGTACCAAGTGTTACACACCAGCACG3' (SEQ ID NO: 12) as a
second strand primer. The PCR products were digested by XhoI+ SalI and cloned into
10 pSPDAUG by XhoI. The resulting plasmid is pNFI/TVE (Fig. 1B).

Example 2. Construction of NF2 to express different foreign peptides.

Assembly of fusion proteins into virus particles greatly enhances the ability to purify the peptides or polypeptides from plants simply by the isolation of the virus
15 (see Example 3). Some of the products (peptides), however, may require separation
from fusion protein (particles) for functional activity. The AIMV coat protein has a
natural trypsin recognition site, which allows cleavage between amino acid 24 and 25
at the N-terminus of the coat protein. When the peptides are fused to NF1, the
cleavage with trypsin will result in a foreign peptide carrying 24 N-terminal amino
20 acids of AIMV CP at their C-terminus. This can be detrimental for the functional
activity of some peptides. To improve this feature of the carrier molecule, a number of
AIMV coat protein mutants were tested. The deletion of 12 N-terminal amino acids of
AIMV coat protein resulted in a mutant molecule, which retains the function of the
wild type and is capable of accommodating the same size peptides as the wild type.
25 The new clone, pSPDATG3 contains all the wild type RNA sequences. It is derived
from pSP65DAUG by creating a KpnI cloning site at position 1226 (RNA3 sequence),
which is 33 nucleotides (11 amino acids) downstream from the original translation
initiation site (AUG). The KpnI site at position 1226 was introduced by PCR using

5'GCGCTCGAGTTCTTCACAAAAGAAAGCTGGTGGGAAAAGGTACCGCTGG
TAAACCT3' (SEQ ID NO: 13) as a first strand primer and

5'ATTAAAAGAGCTCAGACTC3' (SEQ ID NO: 14) as second strand primers. The
PCR product was digested by XhoI+SstI and cloned into pSP65ΔAUG cleaved by

5 XhoI+SstI to replace the original DNA fragment with a mutant one (Fig. 2).

Illustrated in figure 2 is a schematic representation of cloning of foreign
peptides as a translational fusions with mutant AIMV CP (CPATG3) in full-length
RNA3 of AIMV. (A) represents cloning strategy used. Two ellipsoids represent a
foreign peptide with KpnI cloning site. CPATG3 is a mutant AIMV CP where the
10 translation initiation codon (AUG) is mutated to create Xho I site and KpnI at position
1226 for the cloning of foreign peptides. T7P3Sal is 5' portion of AIMV RNA3
containing ORF for P3, 5' non coding regions of RNA3 and subgenomic RNA4. The
Sal I site is created at the position 1192 to mutate the AIMV CP AUG and to clone
into Xho I site at the same position in CPATG3. Additionally T7P3Sal contains T7
15 promoter for in vitro synthesis of infectious RNA3 transcripts. NF2 is full-length
RNA3 containing foreign peptide fused to AIMV CP. (B) pNF2/RSV contains 24
amino acid epitope from RSV G protein. pNF2/Sand contains octerotide sandostatin
and pNF2/TVE contains CP fused with 104 amino acids from colorectal cancer
GA733-2. Amino acid sequences of cloned peptides are shown under each construct.
20 Stem-loop structure indicates the 3' non coding region of RNA3.

To compare the mutant and wild type protein we engineered the fusions with
the following peptides:

a. **pNF2/RSV** (Fig. 2B). DNA sequence coding for antigenic epitope of
RSV G protein was PCR amplified using
25 5'GCGCTCGAGGGTACCATGTCCCTTGTACCCTGCAGCATATGCAGCAACAA
TCCA3' (SEQ ID NO: 15) as a first strand and
5'CGAGGTACCCTCTGGTATTC'TTTTGCAGATAGCCCAGCAGGTTGGATTGT
TGCT3' (SEQ ID NO: 16) as second strand primers. During PCR the KpnI site was

introduced for cloning into NF2. The PCR product was digested by KpnI and ligated into NF2 linearized by KpnI to obtain NF2RSV. pNF2/RSV consists of full-length RNA3 where the antigenic epitope of RSV G protein is fused to the N-terminus of mutant coat protein CPDATG3.

- 5 b. **pNF2/Sand** (Fig. 2B). The sequences encoding sandostatin were PCR amplified using
 5'GCGGGTACCATGTTCTGTTTCTGGAACGTTACTGCTGGTAAACCTAC
 TAAACGT3' (SEQ ID NO: 17) as a first strand and
 5'GCGCTCGAGCATCCCTTAGGGGCATTCATGCA3' (SEQ ID NO: 18) as second
 10 strand primers. The first strand primer contains sequences for both sandostatin (27 nucleotides) and A1MV CP (19 3' nucleotides) for annealing. Thus, the PCR product will contain sequences encoding sandostatin and all the sequences of A1MV CP downstream of the nucleotide 1226 of RNA3. Therefore, the PCR product was
 15 digested by KpnI (newly introduced) Apal (in original A1MV sequences), where the sequences encoding sandostatin were followed by the sequences encoding mutant coat protein CPDATG3 and were cloned into NF2RSV by KpnI Apal to replace an identical region of coat protein together with the fused RSV epitope. The resulting plasmid is pNF2/Sand (Fig. 2B).

- c. **pNF2/TVE** (Fig. 2B). 104 amino acid peptides from colorectal cancer
 20 antigen GA733-2 (Linnenbach et al., 1989, *Proc. Natl. Acad. Sci. USA* 86: 27-31) were fused with mutant A1MV CP. This region has special conformation recognized by colorectal cancer associated antibody 17-1A. The sequences encoding this region (80 amino acids) of GA733-2 were PCR amplified using
 5'GCGCTCGAGGGTACCATGCG ACGGCGACTTTTGCCGCA3' (SEQ ID NO:
 25 19) as a first strand and 5'GTCGACCTGGTACCAGTGTT CACACACCAGCAGC3' (SEQ ID NO: 20) as second strand primers. The PCR products were digested by KpnI and cloned into NF2RSV by Kpn I to replace the RSV epitope fused with the mutant coat protein. The resulting plasmid is pNF2/TVE (Fig. 2B).

Example 3. Construction of RNA3a and RNAM to produce full-length proteins.

A1MV has three genomic RNAs. RNA1 and 2 encode for P1 and P2 proteins
 5 required for the replication of viral RNA. RNA3 encodes for P3 (cell to cell
 movement) and coat protein (long distance movement and encapsidation). Coat
 protein is translated from subgenomic RNA4. RNA4 is synthesised from genomic
 RNA3. P3 and coat protein are required for virus to be fully infectious. Deletion of
 either of these two proteins will limit the infectivity of viruses. Based on the
 10 importance of these two proteins for virus infectivity we introduced mutations into
 RNA3 to create two new molecules (RNA3a and RNA3b, Fig. 2). RNA3a has
 functionally active P3 and is deficient in coat protein production. RNA3b has
 functionally active coat protein and is deficient in P3 production. The functions of coat
 protein and P3 can be complemented from two different molecules (RNA3a and
 15 RNA3b) replacing the wild type RNA3. The RNA3a is NF2 (see Example 2) where
 the wild type coat protein is replaced with mutant coat protein. The mutations were
 introduced to eliminate the A1MV CP translation initiation codon and to create KpnI
 site for subcloning at position 1226. Thus, there is no translation initiation codon for
 coat protein gene in RNA3a. Instead it has XhoI (position 1192 of RNA3), KpnI and
 20 Apal (position 1800 of RNA3) cloning sites for replacing the coat protein sequences
 with the sequences of the desired gene using XhoI-Apal or KpnI-Apal (Fig. 2).

In RNA3b we replaced the translation initiation codon for P3 (ATG) with NheI
 restriction site by PCR using
 5'GCACTCATTCAACATTGCTAGCTTATGTTTTTGTTCACGGAGCTCAAG3'
 25 (SEQ ID NO: 21) as a second strand primer and
 5'CATGCCATTGAMAGGTGACACAATAG3' (SEQ ID NO: 22) as a first strand
 primer. As a template we used T7RNA3. The amplified DNA fragment containing T7
 promoter and 5' 245 nucleotides of RNA3 with mutation was digested by PstI XhoI

and cloned into T7RNA3 cleaved by PstI XhoI to replace the wild type sequences. Thus, RNA3b has XhoI and NdeI restriction sites (Fig. 4) for replacing the open reading frame of P3 with the sequences of the desired gene. This system allows expression of 1 or 2 genes simultaneously. Using this strategy we cloned the following

5 proteins into RNA3a or RNA3b (Fig. 3A or 4A):

Shown in figure 3 is the construction of RNA3a. RNA3 is wild-type genomic RNA of AIMV. The boxes indicate the ORF's for movement protein P3 and for coat protein CP. Stem-loop structure indicates the 3' non coding region of RNA3. (A) The PCR product containing newly introduced KpnI site and mutated AUG codon is

10 cloned into CPDAUG to create CPATG3. CPATG3 is deficient in translation of AIMV CP and has KpnI cloning site at position 1226. Then the CPATG3 is combined with T7P3Sal using Xho I and Sal I sites for cloning as described in Fig. 2 to obtain RNA3a. (B) A3a/GFP- the ORF for AIMV CP is replaced with that of GFP. A3a/17-1ALC contains the ORF for LC of colorectal cancer associated antibody 17-1A.

15 A3a/gp53 contains the glycoprotein from bovine viral diarrhoea virus.

Construction of RNA3b is shown in figure 4 RNA3 is wild-type genomic RNA of AIMV. The boxes indicate the ORF's for movement protein P3 and for coat protein CP. Stem-loop structure indicates the 3' non coding region of RNA3. (A) The PCR product containing T7 promoter and mutated AUG codon of P3 is cloned into

20 T7/A3 the infectious cDNA clone of AIMV RNA3 using Pst I+Xho I restriction sites. The resulting plasmid is RNA3b. RNA3b is deficient in translation of P3 and has Xho I(position 245) NdeI (positions 1082) cloning sites. (B) shows A3b/GFP-in which the ORF for P3 is replaced with that of GFP. Stem-loop structure indicates the 3' non coding region of RNA3.

25 a. GFP (green fluorescent protein) from jellyfish (Fig. 3B) GFP has been used as a marker for expression in different systems. We amplified GFP to introduce KpnI and Apal restriction sites for cloning using

5'GCGGGTACCGTCGACGGCCAGATCGCCATGAGTAAAGGAGAAGAAC3'

(SEQ ID NO: 23) as a first strand and

5'GCGGGCCCATTAATGCGGCCGCTCATTTGTATAGTTCATCC3' (SEQ ID NO: 24) as second strand primers. The amplified PCR product was digested by KpnI+ApaI and cloned into pNF2/RSV cleaved by KpnI+ApaI. The resulting clone pA3a/GFP contains 5'- and 3'-non coding regions of RNA3 and ORF's for P3 and GFP. The open reading frame of the coat protein between nucleotides 1226-1800 was replaced with an open reading frame of GFP.

b. gp53 of Bovine Viral Diarrhea Virus (BVDV, Fig. 3B). gp53 is shown to stimulate virus neutralizing antibodies in vitro and in vivo. gp53 is a major component of diagnostic kits as well as vaccine preparations used to detect and prevent BVDV. We cloned the open reading frame of gp53 from NADI strain of BVDV by RT-PCR into pGEM-T (Promega, Madison, Wisconsin) expression vector using 5'GCGGGCCCATTAATGCGGCCGCTCATTTGTATAGTTCATCC3' (SEQ ID NO: 25) as a second strand and

5'GCAC'CGAGTTACTCACTTGATATGATTTTCATATGGTCT3' (SEQ ID NO: 26) as first strand primers. After sequence confirmation, the ORF of gp53 was cleaved by PaeI+XhoI and cloned into pNF2/RSV digested by KpnI+ApaI using blunt end ligation to create pA3a/gp53 (Fig. 3B).

c. Light chain (LC) of monoclonal antibody 17-1A (Fig. 3B). 17-1A is the monoclonal antibody (Linnenbach et al., 1989, *Proc. Natl. Acad. Sci. USA* 86: 27-31) against colorectal cancer associated with GA733-2 antigen. The gene encoding light chain of 17-1A was amplified using

GCGTTAATTAAGGCCAGATCGGCCATGGGCATCAAGATGGGATCA3' (SEQ ID NO: 27) as a first strand- and

5'GCGTTAATTAAGCGGCCGCC'AACTCATTTCCTGTTGAA3' (SEQ ID NO: 28) as a second-strand primer and cloned into pGEM-T vector. After sequence confirmation the DNA fragment containing ORF of 17-1A LC was cleaved by PaeI and

cloned into pNF2/RSV digested by KpnI+ApaI using blunt end ligation to create pA3a/17-1ALC (Fig. 3B).

- d. GFP was also cloned into RNA3b to replace the ORF for P3 and to create pA3b/GFP (Fig. 4B). The ORF of GFP was cleaved from pGEM-GFP by KpnI+ApaI and cloned into RNA3b cleaved by NheI+NdeI to replace the ORF of P3. The resulting clone pA3b/GFP (Fig. 4B) contains 5'- and 3'-non coding regions of RNA3, AIMV CP and the GFP ORF. The open reading frame of P3 between nucleotides 245-1100 was replaced with an open reading frame of GFP.

Example 4. Production of foreign peptides fused to AIMV CP using transgenic plants expressing the replicase proteins of AIMV.

Rep plants are transgenic tobacco plants expressing replicase proteins (P1 and P2) of AIMV. It has been demonstrated that inoculation of these plants with RNA3 only results in virus infection and systemic movement of RNA3. This shows that P1 and P2 expressed in transgenic plants will complement for virus replicase function.

To test our hypothesis of producing foreign peptides fused to AIMV coat protein using transgenic Rep plants, we inoculated the plants with in vitro transcripts of pNF1/RSV, pNF1/Sand, Pnf1/g24, pNF1/TVE, pNF2/RSV, pNF2/Sand and pNF2/TVE. Tobacco leaves were inoculated with in vitro transcription products of recombinant RNA3. The transcription products of recombinant RNA3 were diluted 1:1 in 30 mM sodium phosphate, pH 7.2, and applied onto expanding tobacco leaves after abrading the leaf surface with carborundum (320 grit; Fisher, Pittsburgh, PA). Inoculation was affected by gentle rubbing to spread the inoculum and further abrade the leaf surface. The recombinant virus was isolated 12-14 days after the inoculum was applied, as described (Yusibov et al., 1997, *Proc. Natl. Acad. Sci. USA* **94**, 5784-5788). Briefly, leaf tissue was ground and the sap separated from cell debris by centrifugation. Virus particles were selectively precipitated using 5% polyethylene

glycol. Then the purified virus was analyzed for the presence of full-length recombinant protein and the peptide of interest using Western analysis.

Western Blot Analysis: Recombinant proteins produced in virus-infected plants were analyzed by Western blot (Yusibov et. al., 1997, *Proc. Natl. Acad. Sci. USA* **94**, 5784-5788). Proteins from purified virus particles were separated electrophoretically on SDS-polyacrylamide gels and electroblotted onto a nylon membrane overnight at 33 mA. After blocking with milk (Kirkegaard & Perry; Gaithersburg, MD), proteins were allowed to react with appropriate antibodies and detected using a Vectastain ABC kit (Vector Laboratories, Burlingame, CA). The recombinant proteins were detected using antibodies specific for: the AIMV coat protein (Agdia, Elkhart, IN); the linear epitope (G5-24) of rabies glycoprotein (Dietzschold et. al., 1990, *J Virology* **64**, 3804-3809); .

Shown in figure 5 is Western analysis and Coomassie staining of NF1/RSV. Proteins were separated by electrophoresis through a 13% SDS-polyacrylamide gel and bound with monoclonal antibodies specific for the AIMV coat protein (A), for the epitope of RSV G protein and stained with Coomassie (C). Wild type AIMV coat protein (24 kD) bound only with antibodies against AIMV coat protein (A) and did not bind with antibodies against fusion peptide (B). The fusion protein NF1/RSV, however, was recognized (A and B) with antibodies specific for both carrier molecule (AIMV CP) and fused peptide (RSV) in total extracts from infected leaves as well in purified virus samples. Total extracts (C-(total)) from noninoculated plants did not react with either of the antibodies. Coomassie staining of NF1/RSV in total extracts and in purified virus samples demonstrates the efficacy of purification procedure.

Shown in figure 6 is Western analysis and Coomassie staining of NF2/RSV. Proteins were separated by electrophoresis through a 13% SDS-polyacrylamide gel and bound with monoclonal antibodies specific for the AIMV coat protein (A), for the epitope of RSV G protein and stained with Coomassie (C). Wild type AIMV coat protein (24 kD) bound only with antibodies against AIMV coat protein (A) and did not

bind with antibodies against fusion peptide (**B**). The fusion protein NF2/RSV, however, was recognized (**A and B**) with antibodies specific for both carrier molecule (AIMV CP) and fused peptide (RSV) in total extracts from infected leaves as well as in purified virus samples. Total extracts (C-(total)) from noninoculated plants did not
5 react with either of the antibodies.

Shown in Figure 7 is Western analysis and Coomassie staining of NF2/Sand. Proteins were separated by electrophoresis through a 13% SDS-polyacrylamide gel and bound with monoclonal antibody specific for the AIMV coat protein (**A**) and stained with Coomassie (**B**). The antibody reacted with AIMV coat protein (24 kD) and with fusion protein NF2/Sand (**A**). The NF2/Sand was recognized (**A and B**) with
10 antibody in total extracts from infected leaves as well in purified virus samples. Total extracts (C-(total)) from noninoculated plants did not react with either of the antibodies. Coomassie staining of NF2/Sand in total extracts and in purified virus samples demonstrates the efficacy of purification procedure.

Shown in figure 8 is Western analysis of NF1/TVE and NF2/TVE accumulation and assembly in infected plants. Proteins were separated by electrophoresis through a 13% SDS-polyacrylamide gel and bound with monoclonal antibodies specific for the AIMV coat protein (**A**) and for colorectal cancer antigen GA733-2 (**B**- NF1/TVE and **C**- NF2/TVE). Wild type AIMV coat protein (24 kD) bound only with antibodies against AIMV coat protein (**A**) and did not bind with
20 antibodies against fusion peptide (**B and C**). The fusion proteins NF1/TVE and NF2/TVE, however, were recognized (**A, B and C**) with antibodies specific for both carrier molecule (AIMV CP) and fused peptide (TVE) in total extracts from infected leaves as well in purified virus samples. Total extracts (C-(total)) from noninoculated
25 plants did not react with either of the antibodies.

a. NF1RSV. Systemically infected leaf tissue was harvested 14-16 days after inoculation for the analysis of NF1RSV accumulation and assembly into particles. Recombinant virus was purified from infected tissue using standard virus

purification procedures (Welter et. al., 1996, *Vaccines: New technologies & applications*. Cambridge Healthtech Institute's; Yusibov et. al, 1997, *Proc. Natl. Acad. Sci. USA* **94**, 5784-5788). Samples for the analysis (Western and Coomassie) were taken from crude extracts of infected tissue and from purified recombinant virus.

- 5 Recombinant protein was detected using monoclonal antibodies for AIMV CP (Fig. 5A) and for the G protein of RSV (Fig. 5B). The monoclonal antibody for AIMV CP reacted with control protein (AIMV CP; Fig. 5A) and recombinant protein of expected size (kD) in crude extracts [NF1RSV(total); Fig. 5A] and in purified virus sample [NF1RSV(purified); Fig. 5A]. The antibodies for G protein recognized only
- 10 recombinant protein in crude extracts (NF1RSV, (total); Fig. 5B) and in purified virus sample (NF1RSV, purified); Fig 5B) and did not react with AIMV CP alone. The proteins in crude extracts from non-inoculated plants did not bind either of the antibodies (C. total); Figs 5A and B). The Figure 5 C is a Coomassie staining of proteins from crud extracts before virus purification (NF1RSV, total and C, total) from
- 15 isolated virus sample which shows the effectiveness of purification procedure.

Example 5. Production of foreign proteins cloned into AIMV RNA3 using transgenic plants expressing the replicase proteins of AIMV.

- Rep plants were inoculated with in vitro transcripts of P3/GFP, P3/gp53,
- 20 P3/17-1ACH and GFP/CP as described in Example 4. Expression of each recombinant protein in upper systemically infected leaves was assessed by Western and Northern analysis.


- a. **P3/17- 1 ACH.** The plants were inoculated with the 1:1 mixture of in vitro transcripts of P3/17-1ACH and RNA3. Within 14 days after inoculation the
- 25 systemically infected tissue was analyzed for the accumulation of AIMV CP and 17-1ACH. Western analysis of the extracts from systemically infected tissue contained both AIMV CP and 17- 1ACH (Fig. 11). The proteins were detected using specific antibodies. Monoclonal antibodies for AIMV CP were used to detect the coat

protein of virus which is indicative of virus replication and movement. The IgG (peroxidase conjugate) was used to detect the 17-1 ACH. This demonstrates that foreign proteins can be produced using AIMV transcomplementation system.

- b. **P3/gp53.** The plants were inoculated with the 1:1 mixture of in vitro transcripts of P3/gp53 and RNA3. 14 days after inoculation locally and systemically infected leaves were analyzed for the accumulation of AIMV CP and P3/gp53. The antibodies for both AIMV CP and P3/gp53 recognized right size proteins. In addition, we purified the virus from systemically infected leaves and used it for isolation of virus RNA. The isolated virus RNA was used for Northern analysis to test if the recombinant RNA P3/gp53 and its subgenomic RNA consisting of gp53 ORF and RNA3 3' noncoding region are encapsidated. The minus sense RNA of gp53 was used as a probe.

Example 6. Construction of TMV vector for the production of foreign proteins by transcomplementing the long distance movement function of virus.

Our hypothesis was to support the systemic movement of defective TMV and produce foreign proteins by transcomplementing this function from another construct.

- Shown in figures 9A-C are schematic representations of the genome of Av (derivative of TMV) and construction of Av/A4 (A), Av/GFP (B), and Av/A4GFP (C): the 126 kD and 183 kD proteins are required for the TMV replication, 30 kD protein is the viral movement protein, and CP is viral coat protein. Arrow under "TMV CP SP" indicates the subgenomic promoter of TMV CP.  is the 3' noncoding region of AIMV. Rz- indicates ribozyme for self-cleavage.

- Av is a construct which is a derivative of TMV. In this construct the translation start codon (ATG) of TMV CP have been replaced with AGA creating a virus defective in production of coat protein. In addition, 42 nucleotides downstream of mutated ATG codon multiple cloning sites Pac I, Pme I, Age I and Xho I were introduced. Av (Fig. 9A) contain full-length TMV defective in coat protein

production. To construct the chimeric Av containing AIMV CP we used pSP65A4 (Loesh-Fries et. al, 1985, *Virology* 146, 177-187) containing full length cDNA of AIMV RNA4. pSP65A4 was digested by EcoR I +Sma I to cleave the DNA fragment containing 5'- and 3'-non coding regions in addition to the open reading frame of
5 AIMV CP. The EcoR I Sma I fragment was blunt ended and cloned into Av linearized by Xho I to create Av/A4 (Fig. 9A). In vitro synthesized transcripts of Av and Av/A4 were used to inoculate the leaves of *Nicotiana benthamiana*, *Nicotiana tabacum* MD609 and *Spinacia oleracea*. Ten days after inoculation samples from locally and systemically infected tissue were analyzed by immunoblot using monoclonal
10 antibodies specific for AIMV coat protein. The coat protein of AIMV was detected both in locally and systemically infected leaves (data not shown). Plants inoculated with in vitro transcripts of Av developed symptoms only on locally inoculated leaves and virus did not move into systemic tissue. Chimeric virus was purified from systemically infected leaves of *Nicotiana benthamiana*, *Nicotiana tabacum* MD609
15 and *Spinacia oleracea*. Polyethyleneglycol precipitated samples were analyzed for the presence of virus particles and AIMV CP. Electron microscopy revealed the presence of rod shaped particles (300 nm in length) similar to that of TMV. Western analysis and Coomassie staining of SDS-PAGE demonstrated the presence of 24 kD protein recognized with monoclonal antibodies for AIMV CP in purified virus particles. This
20 suggests that the genomic RNA of TMV is encapsidated with AIMV CP. We were not able to detect the TMV CP in these samples neither by Western analysis nor by Coomassie staining of SDS-PAGE (data not shown). These experiments demonstrate that the AIMV CP supports long distance movement of TMV and encapsidates TMV RNA into stable, purifiable rod shaped particles.

25 Since the AIMV CP supports the systemic movement of TMV genome, we engineered another Av construct where the GFP was cloned under the control of TMV CP subgenomic promoter. We cleaved GFP (see Example 3) by KpnI+Apal and cloned it into Av linearized by XhoI using blunt end ligation. In a second construct, to

improve the movement functions, we cloned GFP with 5'- and 3'-non coding regions of AIMV CP from A3a/GFP (see Fig. 3 in Example 3). We cleaved the ORF of GFP along with 5'- and 3'-non coding regions of AIMV CP from A3a/GFP by NdeI+SmaI and cloned into Av linearized by XhoI using blunt end ligation to create Av-A3a/GFP (Fig. 9B)

We inoculated *N. benthamiana* plants with the mixture of in vitro synthesized transcripts from Av/A4+Av-A3a/GFP and Av/A4+Av/GFP. Twelve to 15 days after inoculation we analyzed the systemically (upper) infected leaves for the presence of GFP. The control plants inoculated with Av, Av/A4, A4+Av/GFP, and Av-A3a/GFP only had no detectable accumulation of GFP or its messenger RNA in systemically infected leaves. The accumulation of GFP and its messenger RNA in systemically infected leaves was observed using both Western and Northern analysis. The protein was detected using GFP specific antibodies (Clontech). The Northern analysis was performed using 700 bp DNA fragment as a probe. The results of analysis shows that the foreign proteins can be produced in virus infected plants using functional complementation.

Example 7 - Experimental Immunization of Mice with NF1/RSV

Construct Expressing the 25 Amino Acid Antigenic (Protective) Peptide of RSV

G Protein and Challenge with RSV

Eight week old female Swiss-Webster, outbred mice were immunized with 50 µg per dose of recombinant NF1/RSV engineered to express the 25 amino acid antigenic (protective) peptide of RSV G protein. Four immunizations of 0.1 ml were administered intraperitoneally at intervals of 2 weeks each with complete Freund's adjuvant (CFA) at 1:1, volume:volume ratio. An equal quantity of a mixture of wild type AMV was used with CFA as a negative control. Identical peptide (VRS-long), expressed in *Escherichia coli* (*E. coli*) and assembled into inclusion bodies, has been used as a positive control. *E. coli* expressed peptide VRS-long has been demonstrated

to provide complete protection of immunized mice against RSV. Ten to fourteen days after each immunization, serum samples were obtained from individual mice, and RSV-specific antibody titers were assessed. Antigen-specific antibody analysis of serum was performed using an enzyme-linked immunoabsorbant assay (ELISA).

- 5 ELISA plates (Nunc Polysorp, Denmark) were coated with 100 μ l per well of G protein (5 μ g/ml in phosphate-buffered saline) overnight at room temperature (RT; about 25 °C). Coated plates were washed 3 times with PBS-Tween (0.05%) and then blocked with 5% dried milk in PBS at RT for at least one hour. A series of dilutions of sera were added to the plates (30 μ l/well) for 2 to 4 hours at RT. The plates were
- 10 then washed three times with PBS-Tween and peroxidase conjugated secondary antibodies (goat anti-mouse IgG, either whole molecule or gamma chain specific), were added (100 μ l per well) at a final dilution of 1:2000 in PBS, for 1 hour at RT. Plates were the washed 5 times with PBS-Tween a n d TMB substrate was added (100 μ l/well) in phosphate-citrate buffer containing urea, for 30 min at RT in the dark. The
- 15 reaction was stopped with 2M H₂SO₄ (50 μ l per well), and the color change resulting from bound specific antibody measured at 450 nM in an ELISA plate-reader (BioTek, Winooski, VT). The titers are shown in Table 1.

- After the last immunization the mice were internasally challenged with RSV strain A. Then the mice were sacrificed, and the virus load was monitored. While the
- 20 mice immunized with backbone vector AIMV had a high load of virus, the mice immunized with NF1/RSV and VRS-long were protected (Table 1).

Table 1.ELISA titers of sera from mice immunized i.p. with NF1/RSV and challenge infection of these mice with RSV.

Groups of mice	ELISA	CHALLENGE with RSV Protection test: DICT ₅₀ log ₁₀ /g
Mice immunized with NF1/RSV	4,800	\bar{x} : 2.0

Mice immunized with AIMV	300	\bar{x} : 2.9
VRS-long	102400	\bar{x} : 1.7

Example 8 - Engineering and Expression of Full-Length Monoclonal Antibody CO17-1A Associated with Colorectal Cancer

5 Light (LC) and Heavy chains (HC) of CO17-1A were cloned into RNA3 of AIMV. The proteins were expressed using the complementation system described above (Examples 3 and 4). The LC was cloned into RNA3 to replace the coat protein gene and to create A3a17-1ALC (Fig. 3B). The HC was cloned into RNA3 to replace the P3 gene and to obtain A3b17-1AHC.

10 **Plant Inoculation and Protein Extraction:** Leaves of Rep plants were co-inoculated with in vitro transcription products of recombinant A3a17-1ALC and A3b17-1AHC as described (39) after adding 1 vol (v/v) of FES buffer [sodium-pyrophosphate 1% (w/v), malacoid 1% (w/v), celite 1% (w/v), glycine 0.5 M, K₂HPO₄ 0.3 M, pH 8.5, with phosphoric acid. The mixture of in vitro transcription
15 products and FES buffer was applied to tobacco leaves after abrading the leaf surface with carborundum (320 grit; Fisher, Pittsburgh, PA). Inoculation was affected by gentle rubbing to spread the inoculum and further abrade the leaf surface.

Two to three weeks after inoculation, systemically infected leaves were harvested and total soluble protein was extracted by grinding the leaves in 2 vol (w/v)
20 of PE buffer (0.2 M Tris, 5 mM EDT, 0.1% Tween 20). After sedimentation of cell debris in a microtube centrifuge (15 min, 13,000 rpm, 4°C), the supernatant was used for antibody detection by ELISA.

Purification of rAB CO17-1A: Systemically infected leaves were harvested, homogenized in 1 vol (w/v) of phosphate buffer (0.02 M, pH 7.4) and centrifuged (30

min, 14,000 rpm, 4 °C) to remove debris. The supernatant was additionally purified by filtering through a nitrocellulose membrane (0.45 µm pore size). The final extract was applied at 1-ml/min on a 1-ml Sepharose HiTrap/E protein column (Pharmacia, Piscataway, NJ) equilibrated with phosphate buffer. The column was washed with 10
5 vo, (v/v) of phosphate buffer and bound antibody was eluted in 5 vol (v/v) of citrate buffer (0.1 M, pH 4) at 1 ml/min. Fractions of 1, 2, and 3 ml were obtained and analyzed by immunoblot and ELISA.

ELISA: Full-length, assembled rAB was not in the list of references detected by ELISA (Yusibov et. al., 1997, *Proc. Natl. Acad. Sci. USA* 94, 5784-5788). Buffers
10 were prepared as described in Clark et al., 1977. High binding, 96-well ELISA plates (Nunc, F) were coated with Ag GA 733-2 (Dr . D. Herlyn, Wistar Institute, Philadelphia, PA) at a concentration of 1 µg/ml for 1h at 37 °C. Plant extract (antibody) was applied in extraction buffer (0.2 M Tris, 1mM EDTA, 0.1% sodium azide, pH 7.5) and incubated for 2h at 37 C. Bound rAB CO17-1A was detected using
15 an anti-mouse IgG peroxidase conjugate (whole molecule of Fc specific, Sigma. St. Louis, MO).

Expression and Assembly of rAB CO17-1A: Infected leaves of Rep plants were homogenized in extraction buffer, centrifuged to remove cell debris, and directly applied on ELISA plates coated with Ag GA733-2. Plant-produced rAB CO17-1A
20 bound to the Ag was detected using an anti-mouse IgG peroxidase conjugate (whole molecule or Fc). Extracts from plants infected with either A3a17-1ALC or A3b17-1AHC were used as negative controls.

Shown in figure 10 is ELISA analysis of CO17-1A self-assembly in virus-infected plants. At 19 days post-inoculation, systemically infected plant leaves were
25 homogenized in 2 vol (w/v) of extraction buffer, centrifuged to remove cellular debris, and the supernatant (1:2 dilution) was applied on ELISA plates coated with purified Ag GA733-2. Reactivity of the antibody with GA733-2 was detected with anti-mouse IgG peroxidase conjugate, using the whole molecule (A). NI, extract from non-

inoculated control plants; LC and HC , extract from plants expressing only CO17-1A light chain (LC) or heavy chain(HC); LC+HC, extract from plants expressing compete rAb CO17-1A. Data are mean ELISA results (SD) from 5 individual plants.

As shown in Figure 10, the average OD490 nm reading in samples expressing whole antibody (HC+LC) was 2- to 3-fold higher than that of controls (NI, HC, LC) using the whole molecule.

While this invention has been described with a reference to specific embodiments, it will be obvious to those of ordinary skill in the art that variations in these methods and compositions may be used and that it is intended that the invention may be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications encompassed within the spirit and scope of the invention as defined by the claims.

SEQUENCE LISTING

5 <110> Koprowski, Hilary
 Yusibov, Vidadi
 <120> Production Of Biomedical Peptides And Proteins In
 Plants Using Transcomplementation Systems
 <130> JEFF-KOP01
 10 <140> Not Yet Assigned
 <141> Herewith
 <150> US 60/118,867
 15 <151> 1999-02-05
 <160> 28
 <170> PatentIn Ver. 2.0
 20 <210> 1
 <211> 24
 <212> PRT
 <213> respiratory syncytial virus
 25 <300>
 <301> Bastien, N
 Trudel, M
 Simard, C
 30 <302> Protective immune responses induced by the immunization
 of mice with a recombinant bacteriophage displaying an
 epitope of the human respiratory syncytial virus
 <303> Virology
 <304> 234
 35 <306> 118-122
 <307> 1997
 <400> 1
 Met Ser Phe Val Pro Cys Ser Ile Cys Ser Asn Asn Pro Thr Cys Trp
 40 1 5 10 15
 Ala Ile Cys Lys Arg Ile Pro Glu
 20
 45 <210> 2
 <211> 46
 <212> DNA
 <213> Artificial Sequence
 50 <220>

5 <223> Description of Artificial Sequence: Primer
 <400> 2
 gcgctcgagc atcatgtcac cctgcagcat atgcagcaac aatcca 46
 <210> 3
 <211> 41
 <212> DNA
 <213> Artificial Sequence
 10 <220>
 <223> Description of Artificial Sequence: Primer
 <400> 3
 15 cgcgtcgact tgcagatagc ccagcagggtt ggattgttgc t 41
 <210> 4
 <211> 82
 <212> DNA
 20 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Primer
 25 <400> 4
 gcgctcgagg gtaccatgtc cgccgtctac acccgaatta tgatgaacgg aggacgactt 60
 aagcgaccac cagaccagct tg 82
 <210> 5
 30 <211> 93
 <212> DNA
 <213> Artificial Sequence
 <220>
 35 <223> Description of Artificial Sequence: Primer
 <400> 5
 cgaggtagcc tctccacca caagggtgctc atttctgctg gatcggaagt cgtgaagggtt 60
 cacaagctgg tctgggtggc gcttaagtcg tcc 93
 40 <210> 6
 <211> 40
 <212> PRT
 <213> Artificial Sequence
 45 <220>
 <223> Description of Artificial Sequence: A chimera of
 an epitope G5-24 of rabies virus glycoprotein and
 an epitope of rabies virus nucleoprotein
 50

<400> 6
 Met Ser Ala Val Tyr Thr Arg Ile Met Met Asn Gly Gly Arg Leu Lys
 1 5 10 15
 5 Arg Pro Pro Asp Gln Leu Val Asn Leu His Asp Phe Arg Ser Asp Glu
 20 25 30
 Ile Glu His Leu Val Val Glu Glu
 35 40
 10
 <210> 7
 <211> 9
 <212> PRT
 15 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: A peptide to
 suppress the synthesis of human growth hormone
 20
 <400> 7
 Met Phe Cys Phe Trp Lys Thr Cys Thr
 1 5
 25
 <210> 8
 <211> 69
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 <223> Description of Artificial Sequence: Primer
 <400> 8
 35 gcggaattcg ttttatatt taattttctt tcaattactt ccatcatgag ttctttctgt 60
 ttctggaaa 69
 <210> 9
 <211> 35
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> Description of Artificial Sequence: Primer
 45
 <400> 9
 gcgctcgagc gactacacgt ttccagaaa cagaa 35
 50
 <210> 10
 <211> 104

<212> PRT
 <213> Artificial Sequence

 <220>
 5 <223> Description of Artificial Sequence: A peptide from
 colorectal cancer antigen GA733-2

 <300>
 <301> Gronenborn, B
 10 Gardner, R C.
 Schaefer, S
 Shepherd, R J.
 <302> Propagation of foreign DNA in plants using cauliflower
 mosaic virus as vector
 15 <303> Nature
 <304> 294
 <306> 773-776
 <307> 1981

 20 <400> 10
 Met Thr Ala Thr Phe Ala Ala Ala Gln Glu Glu Cys Val Cys Glu Asn
 1 5 10 15

 Tyr Lys Leu Ala Val Asn Cys Phe Val Asn Asn Asn Arg Gln Cys Gln
 25 20 25 30

 Cys Thr Ser Val Gly Ala Gln Asn Thr Val Cys Ser Lys Leu Ala Ala
 35 40 45

 30 Lys Cys Leu Val Met Lys Ala Glu Met Asn Gly Ser Lys Leu Gly Arg
 50 55 60

 Arg Ala Lys Pro Glu Gly Ala Leu Gln Asn Asn Asp Gly Leu Tyr Asp
 65 70 75 80
 35 Pro Asp Cys Asp Ser Gly Leu Phe Lys Ala Lys Gln Cys Asn Gly Thr
 85 90 95

 Ser Thr Cys Trp Cys Val Asn Thr
 40 100

 <210> 11
 <211> 38
 45 <212> DNA
 <213> Artificial Sequence

 <220>
 50 <223> Description of Artificial Sequence: Primer

<400> 11
 gcgctcgagg gtaccatgcg acggcgactt ttgccgca 38

 5 <210> 12
 <211> 34
 <212> DNA
 <213> Artificial Sequence

 10 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 12
 gtcgacctgg taccagtgtt cacacaccag cacg 34

 15 <210> 13
 <211> 56
 <212> DNA
 <213> Artificial Sequence

 20 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 13
 gcgctcgagt tcttcacaaa agaaagctgg tgggaaaagg taccgctggt aaacct 56
 25
 <210> 14
 <211> 19
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 14
 attaaaagag ctcagactc 19
 35
 <210> 15
 <211> 54
 <212> DNA
 40 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 45 <400> 15
 gcgctcgagg gtaccatgtc ctttgtacct tgcagcatat gcagcaacaa tcca 54

 <210> 16
 <211> 54
 50 <212> DNA

<213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Primer
 5
 <400> 16
 cgagggtaccc tctggtattc tttgcagat agcccagcag gttggattgt tgct 54
 <210> 17
 10 <211> 57
 <212> DNA
 <213> Artificial Sequence
 <220>
 15 <223> Description of Artificial Sequence: Primer
 <400> 17
 gcgggtacca tgtctgttt ctggaaaacg tgtactgctg gtaaacctac taaacgt 57
 20 <210> 18
 <211> 32
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> Description of Artificial Sequence: Primer
 <400> 18
 30 gcgctcgagg atcccttagg ggcatcatg ca 32
 <210> 19
 <211> 38
 <212> DNA
 <213> Artificial Sequence
 35 <220>
 <223> Description of Artificial Sequence: Primer
 <400> 19
 40 gcgctcgagg gtaccatgcg acggcgactt ttgccgca 38
 <210> 20
 <211> 34
 <212> DNA
 45 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: Primer
 50 <400> 20

gtcgacctgg taccagtgtt cacacaccag cacg 34

5 <210> 21
 <211> 48
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> Description of Artificial Sequence: Primer

<400> 21
 gcactcattc aacattgcta gcttatgttt ttgtttacgg agctcaag 48

15 <210> 22
 <211> 26
 <212> DNA
 <213> Artificial Sequence

20 <220>
 <223> Description of Artificial Sequence: Primer

<400> 22
 catgccattg amaggtgaca caatag 26

25 <210> 23
 <211> 47
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> Description of Artificial Sequence: Primer

<400> 23
 gcgggtaccg tcgacggcca gatcggccat gagtaaagga gaagaac 47

35 <210> 24
 <211> 41
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> Description of Artificial Sequence: Primer

<400> 24
 gcggggcccat taatgcggcc gctcatttgt atagttcatc c 41

45 <210> 25
 <211> 39
 <212> DNA
 <213> Artificial Sequence

50

<220>
 <223> Description of Artificial Sequence: Primer
 5 <400> 25
 gcattaatta acatcatggg acagggcatt ctgtggcta 39

 <210> 26
 <211> 39
 10 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer
 15 <400> 26
 gcactcgagt tactcacttg atatgattc atatgtct 39

 <210> 27
 20 <211> 45
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer
 25 <400> 27
 gcgttaatta aggccagatc ggccatgggc atcaagatgg gatca 45

 30 <210> 28
 <211> 40
 <212> DNA
 <213> Artificial Sequence

 35 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 28
 40 gcgttaatta agcgccgcc taacctcat tcctgttgaa 40

E CLAIM

1. A method for producing foreign polypeptides in a host plant through functional transcomplementation of a virus, the method comprising:
 - (a) constructing a recombinant viral vector for systemic infection
5 which comprises a recombinant genomic component of the virus comprising a movement protein encoding nucleic acid sequence and a coat protein nucleic acid sequence, and one or more foreign polypeptide-encoding nucleic acid sequences cloned into the recombinant genomic component such that the expression of the recombinant genomic component also results in the expression of at least one of
10 said foreign polypeptide-encoding nucleic acid sequences;
wherein the recombinant viral vector is deficient for the virus replicase function, wherein the host plant is transgenic for expressing replicase genes of a virus to complement the virus replicase function,
 - (b) infecting the host plant at one or more locations with the
15 recombinant viral vector such that the infection of said plant with the recombinant viral vector at one location results in systemic infection in the host plant; and
 - (c) producing said foreign polypeptides in said plant infected with the recombinant viral vector by growing said plant.
- 20 2. The method of claim 1, wherein the virus is an alfalfa mosaic virus, an ilarvirus, a tobamavirus, a cucumovirus or a closterovirus.
3. The method of claim 2, wherein the virus is an alfalfa mosaic virus.
- 25 4. The method of claim 1, wherein the recombinant genomic component includes a full-length RNA3.
5. The method of claim 4, wherein at least one said foreign
30 polypeptide-encoding nucleic acid sequence is cloned into the full-length RNA3 to create N-terminal fusions with the coat protein.

6. The method of claim 1, wherein the foreign polypeptide-encoding nucleic acid sequence comprises a vaccine antigen.

5 7. The method of claim 6, wherein the vaccine antigen is selected from the group consisting of hepatitis B surface antigen, enterotoxin, rabies virus glycoprotein, rabies virus nucleoprotein, Norwalk virus capsid protein, gastrointestinal cancer antigen, G protein of Respiratory Syncytial Virus, Sandostatin or colorectal cancer antigen.

10 8. The method of claim 1, wherein one of two or more foreign polypeptide-encoding nucleic acid sequences is a promoter sequence for in vitro transcription.

15 9. The method of claim 1, wherein said host plant is a dicotyledon or a monocotyledon.

10. A method for producing foreign polypeptides in a host plant using a virus, the method comprising:

20 (a) constructing a recombinant viral vector for systemic infection which comprises a recombinant genomic component of the virus comprising a movement protein encoding nucleic acid sequence and a functional mutant coat protein nucleic acid sequence encoding an amino acid sequence having N-terminal deletions of up to 12 amino acids, and one or more foreign polypeptide-encoding
25 nucleic acid sequences cloned into the recombinant genomic component wherein one of said foreign polypeptide-encoding nucleic acid sequences is fused to the N-terminus of the functional mutant coat protein nucleic acid sequence such that the expression of the recombinant genomic component also results in the expression of fused foreign polypeptide-encoding nucleic acid sequence;

30 (b) providing the host plant for viral infection;

(c) infecting said plant at one or more locations with the recombinant viral vector such that the infection of said plant with the recombinant viral vector at one location results in systemic infection in the host plant; and

(d) producing said foreign polypeptides in said plant infected with the recombinant viral vector by growing said plant.

11. The method of claim 10, wherein the virus is an alfalfa mosaic virus, an ilarvirus, a tobamavirus, a cucumovirus or a closterovirus.

12. The method of claim 11, wherein the virus is an alfalfa mosaic virus.

13. The method of claim 10, wherein the host plant is a non-transgenic plant or a transgenic plant that is transgenic for expressing replicase genes of the virus.

14. The method of claim 13, wherein the amino acid sequence of the functional mutant coat protein has 12 amino acids deleted from the N-terminus.

15. The method of claim 10, wherein the foreign polypeptide-encoding nucleic acid sequence that is fused to the N-terminus of the functional mutant coat protein encodes a vaccine antigen.

16. The method of claim 15, wherein the vaccine antigen is selected from the group consisting of hepatitis B surface antigen, enterotoxin, rabies virus glycoprotein, rabies virus nucleoprotein, Norwalk virus capsid protein, gastrointestinal cancer antigen, G protein of Respiratory Syncytial Virus, Sandostatin or colorectal cancer antigen.

17. The method of claim 10, wherein one of two or more foreign polypeptide-encoding nucleic acid sequences is a promoter sequence for in vitro transcription.

5 18. The method of claim 10, wherein said host plant is a dicotyledon or a monocotyledon.

19. A method for producing foreign polypeptides in a host plant through functional transcomplementation of a virus, the method comprising:

10 (a) constructing a first recombinant viral vector for infection which comprises a recombinant genomic component of the virus comprising the native movement protein encoding nucleic acid sequence and a foreign polypeptide-encoding nucleic acid sequence in place of the native coat protein encoding nucleic acid sequence such that the expression of the recombinant genomic component also
15 results in the expression of the foreign polypeptide-encoding nucleic acid sequence;

(b) constructing a second recombinant viral vector for infection which comprises a recombinant genomic component of the virus comprising the native coat protein encoding nucleic acid sequence and a foreign polypeptide-
20 encoding nucleic acid sequence in place of the native movement protein encoding nucleic acid sequence such that the expression of the recombinant genomic component also results in the expression of the foreign polypeptide-encoding nucleic acid sequence;

(c) infecting said plant at one or more locations with a mixture of
25 the first recombinant viral vector and a second recombinant viral vector such that the infection of said plant with the first and second recombinant viral vectors at one location results in systemic infection in said plant,

wherein the first and second recombinant viral vectors are deficient for the virus replicase function, wherein the host plant is transgenic for expressing
30 replicase genes of a virus to complement the virus replicase function,

wherein the first recombinant viral vector expressing the native movement protein complements the cell-to-cell movement function of the virus and the second recombinant vector expressing the native coat protein complements the long distance transport function of the virus; and

- 5 (d) producing said foreign polypeptides in the host plant infected with the recombinant viral vector by growing the host plant.

20. The method of claim 19, wherein the virus is an alfalfa mosaic virus, an ilarvirus, a tobamavirus, a cucumovirus or a closterovirus.

10

21. The method of claim 20, wherein the virus is an alfalfa mosaic virus.

22. The method of claim 19, wherein the recombinant genomic component includes a full-length RNA3.

15

23. The method of claim 19, wherein the foreign polypeptide-encoding nucleic acid sequence comprises a vaccine antigen.

20 24. The method of claim 23, wherein the vaccine antigen is selected from the group consisting of hepatitis B surface antigen, enterotoxin, rabies virus glycoprotein, rabies virus nucleoprotein, Norwalk virus capsid protein, gastrointestinal cancer antigen, G protein of Respiratory Syncytial Virus, Sandostatin or colorectal cancer antigen.

25

25. The method of claim 19, wherein said host plant is a dicotyledon or a monocotyledon.

26. A method for producing foreign polypeptides in a host plant through functional transcomplementation of a chimeric virus, the method comprising:

30

- (a) constructing a recombinant viral vector for systemic infection which comprises a recombinant genomic component of a first class of virus comprising a movement protein encoding nucleic acid sequence of the first class of virus, a full-length coat protein nucleic acid sequence of a second class of virus in place of the native coat protein nucleic acid sequence of the first class of virus and one or more foreign polypeptide-encoding nucleic acid sequences cloned into the recombinant genomic component such that the expression of the recombinant genomic component also results in the expression of at least one of said foreign polypeptide-encoding nucleic acid sequences;
- (b) infecting the host plant at one or more locations with the recombinant viral vector such that the infection of the host plant with the recombinant viral vector at one location results in systemic infection in the host plant,
- wherein the recombinant viral vector expressing the native movement protein of the first class of virus complements the cell-to-cell movement function of the chimeric virus and that expressing the full-length coat protein nucleic acid sequence of the second class of virus complements the long distance transport function of the chimeric virus; and
- (c) producing said foreign polypeptides in the host plant infected with the recombinant viral vector by growing the host plant.

27. The method of claim 26, wherein the class of virus is a tobamavirus, a cucumovirus or a closterovirus.

28. The method of claim 26, wherein the different class of virus is an alfalfa mosaic virus or ilarvirus.

29. The method of claim 26, wherein at least one of said foreign polypeptide-encoding nucleic acid sequences encodes a vaccine antigen.

30. The method of claim 29, wherein the vaccine antigen is selected from the group consisting of hepatitis B surface antigen, enterotoxin, rabies virus glycoprotein, rabies virus nucleoprotein, Norwalk virus capsid protein, gastrointestinal cancer antigen, G protein of Respiratory Syncytial Virus,
5 Sandostatin or colorectal cancer antigen.

31. The method of claim 26, wherein the plant is a dicotyledon or a monocotyledon.

10 32. A method for producing foreign polypeptides in a host plant through functional complementation of a chimeric virus, the method comprising:

(a) constructing a first recombinant viral vector for systemic infection which comprises a recombinant genomic component of a first class of virus comprising a movement protein encoding nucleic acid sequence of the first
15 class of virus, a non-functional coat protein nucleic acid sequence of the first class of virus, a full-length coat protein nucleic acid sequence of a second class of virus inserted into the non-functional coat protein nucleic acid sequence of the first class of virus;

(b) constructing a second recombinant viral vector for infection
20 which comprises a recombinant genomic component of the first class of virus comprising a movement protein encoding nucleic acid sequence of the first class of virus, a non-functional coat protein nucleic acid sequence of the class of virus, and a first foreign polypeptide-encoding nucleic acid sequences cloned into the recombinant genomic component such that the expression of the recombinant
25 genomic component also results in the expression of the first foreign polypeptide-encoding nucleic acid sequence;

(c) infecting the host plant at one or more locations with the first and second recombinant viral vectors such that the infection of the host plant with said recombinant viral vectors at one location results in systemic infection in the
30 host plant.

wherein the first and second recombinant viral vectors expressing the native movement protein of the first class of virus provide the cell-to-cell movement function of the chimeric virus and the first recombinant viral vector expressing the full-length coat protein nucleic acid sequence of the second class of virus provides the long distance transport function of the chimeric virus; and

(d) producing said foreign polypeptides in the host plant infected with the recombinant viral vector by growing the host plant.

33. The method of claim 32, further comprising constructing a third recombinant viral vector for systemic infection which comprises a recombinant genomic component of the first class of virus comprising a movement protein encoding nucleic acid sequence of the first class of virus, a non-functional coat protein nucleic acid sequence of the first class of virus, and a second foreign polypeptide-encoding nucleic acid sequence together with the 5' and 3' non-coding sequences of the full-length coat protein nucleic acid sequence of the second class of virus inserted into the non-functional coat protein nucleic acid sequence of the first class of virus such that the expression of the recombinant genomic component also results in the expression of the foreign polypeptide-encoding nucleic acid sequence.

20

34. The method of claim 32, wherein the class of virus is, a tobamavirus, a cucumovirus or a closterovirus.

35. The method of claim 32, wherein the different class of virus is an alfalfa mosaic virus or ilarvirus.

25

36. The method of claim 32, wherein at least one of said foreign polypeptide-encoding nucleic acid sequences encodes a vaccine antigen.

37. The method of claim 36, wherein the vaccine antigen is selected from the group consisting of hepatitis B surface antigen, enterotoxin, rabies virus glycoprotein, rabies virus nucleoprotein, Norwalk virus capsid protein, gastrointestinal cancer antigen, G protein of Respiratory Syncytial Virus,
5 Sandostatin or colorectal cancer antigen.

38. The method of claim 32, wherein the host plant is a dicotyledon or a monocotyledon.

10 39. A composition comprising a recombinant viral vector capable of systemic infection for producing foreign polypeptides in a viral replicase expressing transgenic host plant that complements the viral replicase function which composition comprises a recombinant genomic component of the virus comprising a movement protein encoding nucleic acid sequence and a coat protein
15 nucleic acid sequence, and one or more foreign polypeptide-encoding nucleic acid sequences cloned into the recombinant genomic component such that the expression of the recombinant genomic component also results in the expression of at least one said foreign polypeptide-encoding nucleic acid sequence.

20 40. A composition comprising a recombinant chimeric viral vector capable of systemic infection for producing foreign polypeptides in a host plant which comprises a recombinant genomic component of a first class of virus comprising a movement protein encoding nucleic acid sequence of the first class of virus, a coat protein nucleic acid sequence of a second class of virus in place of the
25 native coat protein nucleic acid sequence of the first class of virus and one or more foreign polypeptide-encoding nucleic acid sequences cloned into the recombinant genomic component of the first class of virus such that the expression of the recombinant genomic component also results in the expression of at least one said foreign polypeptide-encoding nucleic acid sequence.

30

41. A composition comprising a mixture of recombinant viral vectors capable of systemic infection for producing foreign polypeptides in a host plant, the mixture comprising:

5 (a) a first recombinant viral vector which comprises a recombinant genomic component of the virus comprising a movement protein encoding nucleic acid sequence and a foreign polypeptide-encoding nucleic acid sequence in place of the native coat protein encoding nucleic acid sequence such that the expression of the recombinant genomic component also results in the expression of the foreign polypeptide-encoding nucleic acid sequence; and

10 (b) a second recombinant viral vector which comprises a recombinant genomic component of the virus comprising the native coat protein encoding nucleic acid sequence and a foreign polypeptide-encoding nucleic acid sequence in place of the native movement protein encoding nucleic acid sequence such that the expression of the recombinant genomic component also results in the
15 expression of the foreign polypeptide-encoding nucleic acid sequence.

42. A composition comprising a plant tissue having foreign polypeptides produced according to the method of claim 1.

20 43. A composition comprising a plant tissue having foreign polypeptides produced according to the method of claim 10.

44. A composition comprising a plant tissue having foreign polypeptides produced according to the method of claim 19.

25 45. A method for eliciting an immunological response in a mammal comprising the step of: administering to the mammal an amount of a polypeptide containing plant or plant tissue thereof produced as in claim 6 to induce an immunological response to said polypeptide in said individual.

30

46. A recombinant in vitro transcript of an alfalfa mosaic virus for producing foreign polypeptides in a host plant, the recombinant in vitro transcript comprising:

- 5 (a) a movement protein encoding nucleic acid sequence;
(b) a coat protein encoding nucleic acid sequence; and
(c) a foreign polypeptide-encodingprotein encoding nucleic acid sequence inserted into the recombinant genomic component such that the expression of the movement protein encoding nucleic acid sequence and the coat protein encoding nucleic acid sequence also results in the expression of the foreign
10 polypeptide-encoding nucleic acid sequence,
wherein the host plant is a transgenic plant that is transgenic for expressing replicase genes of said virus, wherein the transgenic plant expressing replicase genes transcomplements the replicase function of said virus.

47. The recombinant in vitro transcript of claim 46, wherein the foreign
15 polypeptide-encodingprotein encoding nucleic acid sequence is fused to the N-terminus of the coat protein.

1/17




Figure 1A

2/17




Figure 1B

4/17




Figure 2B

3/17




Figure 2A

5/17




Figure 3A




Figure 3B

7/17




Figure 4A

8/17




Figure 4B

9/17

Western Analysis and Coomassie Staining of NF1/RSV




Figure 5

10/17

Western Analysis and Coomassie Staining of NF2/RSV




Figure 6

11/17

Western Analysis and Coomassie Staining of NF2/Sand





Figure 7

Western Analysis of accumulation of NF1/TVE and NF2/TVE



A - AIMV CP specific
antibodies

B and C -GA733-2
specific antibodies

Figure 8

13/17




Figure 9A

14/17




Figure 9B

15/17




Figure 9C

16/17

Detection of Full-Length CO17-1A Produced in Plants by ELISA




Figure 10

17/17

Western Analysis of P3/17-ACH Expression in Plants




Figure 11

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/02971

A. CLASSIFICATION OF SUBJECT MATTER		
IPC(7) : C12N 5/04, 15/63, 15/82, 15/83, 15/90; G01N 33/556		
US CL : Please See Extra Sheet.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
U.S. : 435/ 69.1, 320.1, 410, 419, 468; 436/501; 800/278, 288, 295, 298, 317, 320		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
West, Agricola, Biosis, Caplus		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98/08375 A1 (THOMAS JEFFERSON UNIVERSITY). 05 March 1998. page 19, line 20- page 21, line 25, page 23, line 9- page 28, line 3, page 32, line 3 -page 34, line 11, page 36, line 3 -page 38, line 15.	1-47
Y	MODELSKA et al. Immunization Against Rabies With Plant-derived Antigen. Proc. Natl. Acad. Sci., USA. March 1998. Vol. 95. pages 2481-2485, see whole document.	6, 7, 15, 16, 23, 24, 29, 30, 36, 37, 45
A, P	SPITSIN et al. Expression Of Alfalfa Mosaic Virus Coat Protein In Tobacco Mosaic Virus (TMV) Deficient In The Production Of Its Native Coat Protein Supports Long-Distance Movement Of A Chimeric TMV. Proc. Natl. Acad. Sci., USA. March 1999. Vol. 96. pages 2549-2533, see whole document.	1-47
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
08 MAY 2000		27 JUN 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231		Authorized officer
Facsimile No. (703) 305-3230		ASHWIN MEHTA
		Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/02971

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,866,785 A (DONSON et al) 02 February 1999, col. 5, line 35 -col. 7, line 2, col. 19, line 50 -col. 25, line 5, claims.	1-47

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/02971

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☒ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/02971

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/ 69.1, 320.1, 410, 419, 468; 436/501; 800/278, 288, 295, 298, 317, 320

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-25, 39, 41-44, 46, & 47, drawn to a first method for producing foreign polypeptides in a host plant through functional transcomplementation of a virus, comprising constructing a recombinant viral vector comprising the movement protein and coat protein encoding nucleic acid sequences, and one or more foreign polypeptide-encoding sequences, and a first product, a composition comprising the movement, coat protein, and foreign polypeptide-encoding sequences.

Group II, claim(s) 26-38 & 40, drawn to a second method, for producing foreign polypeptides, comprising a movement protein encoding nucleic acid sequence from a first class of virus, a coat protein encoding sequence from a second class of virus, and a foreign polypeptide encoding sequence, and a second product, comprising a composition of said sequences.

Group III, claim(s) 45, drawn to a third method for eliciting an immunological response in mammals.

The inventions listed as Groups I-III do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the method of Group I comprises using movement and coat protein encoding nucleotide sequences from the same virus, which is not shared with Group II. The method of Group II does not share the two classes of viruses with the methods of the other groups. Group III does not share the eliciting of the mammalian immune response with the other groups.